

1 **CYTOTOXICITY MEDIATION OF CELLS EVIDENCING SURFACE**

2 **EXPRESSION OF CD44**

3
4 Reference to Related Applications:

5 This application is a continuation-in-part of application S.N. 10/603,000, filed June
6 23, 2003, which is a continuation-in-part of application S.N. 09/727,361, filed November
7 29, 2000, which is a continuation-in-part of application S.N. 09/415,278, filed October 8,
8 1999, now U.S. Patent 6,180,357 B1, the contents of each of which are herein incorporated
9 by reference.

10 Field Of The Invention:

11 This invention relates to the diagnosis and treatment of cancerous diseases,
12 particularly to the mediation of cytotoxicity of tumor cells; and most particularly to the use
13 of cancerous disease modifying antibodies (CDMAB), optionally in combination with one
14 or more chemotherapeutic agents, as a means for initiating the cytotoxic response. The
15 invention further relates to binding assays which utilize the CDMAB of the instant
16 invention.

17 Background Of The Invention:

18 Raising monoclonal antibodies against human white blood cells led to the
19 discovery of the CD44 antigen; a single chain hyaluronic acid (HA) binding glycoprotein
20 expressed on a wide variety of normal tissue and on all types of hematopoietic cells. It
21 was originally associated with lymphocyte activation and homing. Currently, its putative
22 physiological role also includes activation of inflammatory genes, modulation of cell cycle,
23 induction of cell proliferation, induction of differentiation and development, induction of
24 cytoskeletal reorganization and cell migration and cell survival/resistance to apoptosis.

McHale & Slavin, P.A.

1 In humans, the single gene copy of CD44 is located on the short arm of
2 chromosome 11, 11p13. The gene contains 19 exons; the first 5 are constant, the next 9
3 are variant, the following 3 are constant and the final 2 are variant. Differential splicing
4 can lead to over 1000 different isoforms. However, currently only several dozen naturally
5 occurring variants have been identified.

6 The CD44 standard glycoprotein consists of a N-terminal extracellular (including a
7 20 a.a. leader sequence, and a membrane proximal region (85 a.a.)) domain (270 a.a.), a
8 transmembrane region (21 a.a.) and a cytoplasmic tail (72 a.a.). The extracellular region
9 also contains a link module at the N-terminus. This region is 92 a.a. in length and shows
10 homology to other HA binding link proteins. There is high homology between the mouse
11 and human forms of CD44. The variant forms of the protein are inserted to the carboxy
12 terminus of exon 5 and are located extracellularly when expressed.

13 A serum soluble form of CD44 also occurs naturally and can arise from either a
14 stop codon (within the variable region) or from proteolytic activity. Activation of cells
15 from a variety of stimuli including TNF- α results in shedding of the CD44 receptor.
16 Shedding of the receptor has also been seen with tumor cells and can result in an increase
17 in the human serum concentration of CD44 by up to 10-fold. High CD44 serum
18 concentration suggests malignancy (ovarian cancer being the exception).

19 The standard form of CD44 exists with a molecular weight of approximately 37
20 kD. Post-translational modifications increase the molecular weight to 80-90 kD. These
21 modifications include amino terminus extracellular domain N-linked glycosylations at
22 asparagine residues, O-linked glycosylations at serine/threonine residues at the carboxy
23 terminus of the extracellular domain and glycosaminoglycan additions. Splice variants can
24 range in size from 80-250 kD.

1 HA, a polysaccharide located on the extracellular matrix (ECM) in mammals, is
2 thought to be the primary CD44 ligand. However, CD44 has also been found to bind such
3 proteins as collagen, fibronectin, laminin etc. There appears to be a correlation between
4 HA binding and glycosylation. Inactive CD44 (does not bind HA) has the highest levels of
5 glycosylation, active CD44 (binding HA) the lowest while inducible CD44 (does not or
6 weakly binds HA unless activated by cytokines, monoclonal antibodies, growth factors,
7 etc.) has glycosylation levels somewhere in between the active and inactive forms.

8 CD44 can mediate some of its functions through signal transduction pathways that
9 depend on the interaction of the cell, stimulus and the environment. Some of these
10 pathways include the NF κ B signaling cascade (involved in the inflammatory response), the
11 Ras-MAPK signal transduction pathway (involved with activating cell cycling and
12 proliferation), the Rho family of proteins (involved with cytoskeleton reorganization and
13 cell migration) and the PI3-K-related signaling pathway (related to cell survival). All of
14 the above-mentioned functions are closely associated with tumor disease initiation and
15 progression. CD44 has also been implicated in playing a role in cancer through a variety of
16 additional mechanisms. These include the presentation of growth factors, chemokines and
17 cytokines by cell surface proteoglycans present on the cell surface of CD44 to receptors
18 involved in malignancy. Also, the intracellular degradation of HA by lysosomal
19 hyaluronidases after internalization of the CD44-HA complex can potentially increase the
20 likelihood of tumor invasiveness and induction of angiogenesis through the ECM. In
21 addition, the transmission of survival or apoptotic signals has been shown to occur through
22 either the standard or variable CD44 receptor. CD44 has also been suggested to be
23 involved in cell differentiation and migration. Many, if not all, of these mechanisms are

environment and cell dependent and several give rise to variable findings. Therefore, more research is required before any conclusions can be drawn.

In order to validate a potential functional role of CD44 in cancer, expression studies of CD44 were undertaken to determine if differential expression of the receptor correlates with disease progression. However, inconsistent findings were observed in a majority of tumor types and this is probably due to a combination of reagents, technique, pathological scoring and cell type differences between researchers. Renal cell carcinoma and non-Hodgkin's lymphoma appear to be the exception in that patients with high CD44 expressing tumors consistently had shorter survival times than their low or non-CD44 expressing counterparts.

Due to its association with cancer, CD44 has been the target of the development of anti-cancer therapeutics. There is still controversy as to whether the standard or the variant forms of CD44 are required for tumor progression. There is *in vivo* animal data to support both views and again it may be tumor type and even cell type dependent. Different therapeutic approaches have included injection of soluble CD44 proteins, hyaluronan synthase cDNA, hyaluronidase, the use of CD44 antisense and CD44 specific antibodies. Each approach has led to some degree of success thereby providing support for anti-CD44 cancer therapeutics.

Both variant and standard CD44 specific monoclonal antibodies have been generated experimentally but for the most part these antibodies have no intrinsic biological activity, rather they bind specifically to the type of CD44 they recognize. However, there are some that are either active *in vitro* or *in vivo* but generally not both. Several anti-CD44 antibodies have been shown to mediate cellular events. For example the murine antibody A3D8, directed against human erythrocyte Lutheran antigen CD44 standard form, was

1 shown to enhance CD2 (9-1 antibody) and CD3 (OKT3 antibody) mediated T cell
2 activation; another anti-CD44 antibody had similar effects. A3D8 also induced IL-1 release
3 from monocytes and IL-2 release from T lymphocytes. Interestingly, the use of A3D8 in
4 conjunction with drugs such as daunorubicin, mitoxantrone and etoposide inhibited
5 apoptosis induction in HL60 and NB4 AML cells by abrogating the generation of the
6 second messenger ceramide. The J173 antibody, which does not have intrinsic activity and
7 is directed against a similar epitope of CD44s, did not inhibit drug-induced apoptosis. The
8 NIH44-1 antibody, directed against an 85-110 KD and 200 KD form of CD44, augmented
9 T-cell proliferation through a pathway the authors speculated as either cross-linking or
10 aggregation of CD44. Taken together, there is no evidence that antibodies such as these are
11 suitable for use as cancer therapeutics since they either are not directed against cancer (e.g.
12 activate lymphocytes), induce cell proliferation, or when used with cytotoxic agents
13 inhibited drug-induced death of cancer cells.

14 Several anti-CD44 antibodies have been described which demonstrate anti-tumor
15 effects *in vivo*. The antibody 1.1ASML, a mouse anti-rat IgG1 directed to the v6 variant of
16 CD44, has been shown to decrease the lymph node and lung metastases of the rat
17 pancreatic adenocarcinoma BSp73ASML. Survival of the treated animals was
18 concomitantly increased. The antibody was only effective if administered before lymph
19 node colonization, and was postulated to interfere with cell proliferation in the lymph
20 node. There was no direct cytotoxicity of the antibody on the tumor cells *in vitro*, and the
21 antibody did not enhance complement-mediated cytotoxicity, or immune effector cell
22 function. Utility of the antibody against human cells was not described.

23 Breyer et al. described the use of a commercially-available antibody to CD44s to
24 disrupt the progression of an orthotopically-implanted rat glioblastoma. The rat

glioblastoma cell line C6 was implanted in the frontal lobe, and after 1 week, the rats were given 3 treatments with antibody by intracerebral injection. Treated rats demonstrated decreased tumor growth, and higher body weight than buffer or isotype-control treated rats. The antibody was able to inhibit adhesion of cells *in vitro* to coverslips coated with extracellular matrix components, but did not have any direct cytotoxic effects on cells. This antibody was not tested against human cells.

A study was carried out which compared the efficacy of an antibody to CD44s (IM-7.8.1) to an antibody to CD44v10 (K926). The highly metastatic murine melanoma line B16F10, which expresses both CD44 isoforms, was implanted i.v. into mice. After 2 days, antibodies were given every third day for the duration of the study. Both antibodies caused a significant reduction of greater than 50% in the number of lung metastases; there was no significant difference in efficacy between the two antibodies. The antibody did not affect proliferation *in vitro*, and the authors, Zawadzki *et al*, speculated that the inhibition of tumor growth was due to the antibody blocking the interaction of CD44 with its ligand. In another study using IM-7.8.1, Zahalka *et al* demonstrated that the antibody and its F(ab')₂ fragment were able to block the lymph node infiltration by the murine T-cell lymphoma LB. This conferred a significant survival benefit to the mice. Wallach-Dayana *et al* showed that transfection of LB-TRs murine lymphoma, which does not spontaneously form tumors, with CD44v4-v10 conferred the ability to form tumors. IM-7.8.1 administration decreased tumor size of the implanted transfected cells in comparison to the isotype control antibody. None of these studies demonstrated human utility for this antibody.

GKW.A3, a mouse IgG2a, is specific for human CD44 and prevents the formation and metastases of a human melanoma xenograft in SCID mice. The antibody was mixed with the metastatic human cell line SMMU-2, and then injected subcutaneously.

1 Treatments were continued for the following 3 weeks. After 4 weeks, only 1 of 10 mice
2 developed a tumor at the injection site, compared to 100% of untreated animals. F(ab')₂
3 fragments of the antibody demonstrated the same inhibition of tumor formation, suggesting
4 that the mechanism of action was not dependent on complement or antibody-dependent
5 cellular cytotoxicity. If the tumor cells were injected one week prior to the first antibody
6 injection, 80% of the animals developed tumors at the primary site. However, it was noted
7 that the survival time was still significantly increased. Although the delayed antibody
8 administration had no effect on the primary tumor formation, it completely prevented the
9 metastases to the lung, kidney, adrenal gland, liver and peritoneum that were present in the
10 untreated animals. This antibody does not have any direct cytotoxicity on the cell line *in*
11 *vitro* or interfere with proliferation of SMMU-2 cells, and appears to have its major effect
12 on tumor formation by affecting metastasis or growth. One notable feature of this antibody
13 was that it recognized all isoforms of CD 44, which suggests limited possibilities for
14 therapeutic use.

15 Strobel *et al* describe the use of an anti-CD44 antibody (clone 515) to inhibit the
16 peritoneal implantation of human ovarian cancer cells in a mouse xenograft model. The
17 human ovarian cell line 36M2 was implanted i.p. into mice in the presence of the anti-
18 CD44 antibody or control antibody, and then treatments were administered over the next
19 20 days. After 5 weeks, there were significantly fewer nodules in the peritoneal cavity in
20 the antibody treated group. The nodules from both the anti-CD44 and control treated
21 groups were the same size, suggesting that once the cells had implanted, the antibody had
22 no effect on tumor growth. When cells were implanted subcutaneously, there was also no
23 effect on tumor growth, indicating that the antibody itself did not have an anti-proliferative
24 or cytotoxic effect. In addition, there was no effect of the antibody on cell growth *in vitro*.

1 VFF-18, also designated as BIWA 1, is a high-affinity antibody to the v6 variant of
2 CD44 specific for the 360-370 region of the polypeptide. This antibody has been used as a
3 ^{99m}Techneium-labelled conjugate in a Phase 1 clinical trial in 12 patients. The antibody
4 was tested for safety and targeting potential in patients with squamous cell carcinoma of
5 the head and neck. Forty hours after injection, 14% of the injected dose was taken up by
6 the tumor, with minimal accumulation in other organs including the kidney, spleen and
7 bone marrow. The highly selective tumor binding suggests a role for this antibody in
8 radioimmunotherapy, although the exceptionally high affinity of this antibody prevented
9 penetration into the deeper layers of the tumor. Further limiting the application of BIWA 1
10 is the immunogenicity of the murine antibody (11 of 12 patients developed human anti-
11 mouse antibodies (HAMA)), heterogenous accumulation throughout the tumor and
12 formation of antibody-soluble CD44 complexes. WO 02/094879 discloses a humanized
13 version of VFF-18 designed to overcome the HAMA response, designated BIWA 4.
14 BIWA 4 was found to have a significantly lower antigen binding affinity than the parent
15 VFF 18 antibody. Surprisingly, the lower affinity BIWA 4 antibody had superior tumor
16 uptake characteristics than the higher affinity BIWA 8 humanized VFF-18 antibody. Both
17 ^{99m}Techneium-labelled and ¹⁸⁶Rhenium-labelled BIWA 4 antibody were assessed in a 33
18 patient Phase 1 clinical trial to determine safety, tolerability, tumor accumulation and
19 maximum tolerated dose, in the case of ¹⁸⁶Re-labelled BIWA 4. There appeared to be
20 tumor related uptake of ^{99m}Tc-labelled BIWA 4. There were no tumor responses seen with
21 all doses of ¹⁸⁶Re-labelled BIWA 4, although a number had stable disease; the dose
22 limiting toxicity occurred at 60 mCi/m². There was a 50-65% rate of adverse events with
23 12 of 33 patients deemed to have serious adverse events (thrombocytopenia, leucopenia
24 and fever) and of those 6, all treated with ¹⁸⁶Re-labelled BIWA 4, died in the course of

1 treatment or follow-up due to disease progression. Two patients developed human anti-
2 human antibodies (HABA). A Phase 1 dose escalation trial of ^{186}Re -labelled BIWA 4 was
3 carried out in 20 patients. Oral mucositis and dose-limiting thrombocytopenia and
4 leucocytopenia were observed; one patient developed a HABA response. Stable disease
5 was seen in 5 patients treated at the highest dose of 60 mCi/m^2 . Although deemed to be
6 acceptable in both safety and tolerability for the efficacy achieved these studies have
7 higher rates of adverse events compared to other non-radioisotope conjugated biological
8 therapies in clinical studies. U.S. Patent Application US 2003/0103985 discloses a
9 humanized version of VFF-18 conjugated to a maytansinoid, designated BIWI 1, for use in
10 tumor therapy. A humanized VFF 18 antibody, BIWA 4, when conjugated to a toxin, i.e.
11 BIWI 1, was found to have significant anti-tumor effects in mouse models of human
12 epidermoid carcinoma of the vulva, squamous cell carcinoma of the pharynx or breast
13 carcinoma. The unconjugated version, BIWA 4, did not have anti-tumor effects and the
14 conjugated version, BIWI 1, has no evidence of safety or efficacy in humans.

15 Mab U36 is a murine IgG1 antibody generated by UM-SCC-22B human
16 hypopharyngeal carcinoma cell immunization and selection for cancer and tissue
17 specificity. Antigen characterization through cDNA cloning and sequence analysis
18 identified the v6 domain of keratinocyte-specific CD44 splice variant epican as the target
19 of Mab U36. Immunohistochemistry studies show the epitope to be restricted to the cell
20 membrane. Furthermore, Mab U36 labeled 94% of the head and neck squamous cell
21 carcinomas (HNSCC) strongly, and within these tumors there was uniformity in cell
22 staining. A 10 patient $^{99\text{m}}\text{Tc}$ -labelled Mab U36 study showed selective accumulation of the
23 antibody to HNSCC cancers ($20.4 \pm 12.4 \%$ injected dose/kg at 2 days); no adverse effects
24 were reported but two patients developed HAMA. In a study of radio-iodinated murine

1 Mab U36 there were 3 cases of HAMA in 18 patients and selective homogenous uptake in
2 HNSCC. In order to decrease the antigenicity of Mab U36 and decrease the rate of HAMA
3 a chimeric antibody was constructed. Neither the chimeric nor the original murine Mab
4 U36 has ADCC activity. There is no evidence of native functional activity of Mab U36.
5 ¹⁸⁶Re-labelled chimeric Mab U36 was used to determine the utility of Mab U36 as a
6 therapeutic agent. In this Phase 1 escalating dose trial 13 patients received a scouting dose
7 of ^{99m}Tc-labelled chimeric Mab U36 followed by ¹⁸⁶Re-labelled chimeric Mab U36. There
8 were no acute adverse events reported but following treatment dose limiting myelotoxicity
9 (1.5 GBq/m²) in 2 of 3 patients, and thrombocytopenia in one patient treated with the
10 maximum tolerated dose (1.0 GBq/m²) were observed. Although there were some effects
11 on tumor size these effects did not fulfill the criteria for objective responses to treatment. A
12 further study of ¹⁸⁶Re-labelled chimeric Mab U36 employed a strategy of using
13 granulocyte colony-stimulating factor stimulated whole blood reinfusion to double the
14 maximum-tolerated activity to 2.8 Gy. In this study of nine patients with various tumors of
15 the head and neck 3 required transfusions for drug related anemia. Other toxicity includes
16 grade 3 myelotoxicity, and grade 2 mucositis. No objective tumor responses were reported
17 although stable disease was achieved for 3-5 months in five patients. Thus, it can be seen
18 that although Mab U36 is a highly specific antibody the disadvantage of requiring a
19 radioimmunoconjugate to achieve anti-cancer effects limits its usefulness because of the
20 toxicity associated with the therapy in relation to the clinical effects achieved.

21 To summarize, a CD44v6 (1.1ASML) and CD44v10 (K926) monoclonal antibody
22 have been shown to reduce metastatic activity in rats injected with a metastatic pancreatic
23 adenocarcinoma or mice injected with a malignant melanoma respectively. Another anti-
24 CD44v6 antibody (VFF-18 and its derivatives), only when conjugated to a maytansinoid or

1 a radioisotope, has been shown to have anti-tumor effects. Anti-standard CD44
2 monoclonal antibodies have also been shown to suppress intracerebral progression by rat
3 glioblastoma (anti-CD44s), lymph node invasion by mouse T cell lymphoma (IM-7.8.1) as
4 well as inhibit implantation of a human ovarian cancer cell line in nude mice (clone 515),
5 lung metastasis of a mouse melanoma cell line (IM-7.8.1) and metastasis of a human
6 melanoma cell line in SCID mice (GKW.A3). The radioisotope conjugated Mab U36 anti-
7 CD44v6 antibody and its derivatives had anti-tumor activity in clinical trials that were
8 accompanied by significant toxicity. These results, though they are encouraging and
9 support the development of anti-CD44 monoclonal antibodies as potential cancer
10 therapeutics, demonstrate limited effectiveness, safety, or applicability to human cancers.

11 Thus, if an antibody composition were isolated which mediated cancerous cell
12 cytotoxicity, as a function of its attraction to cell surface expression of CD44 on said cells,
13 a valuable diagnostic and therapeutic procedure would be realized.

14

15 Prior Patents:

16 U.S. Patent No. 5,750,102 discloses a process wherein cells from a patient's tumor
17 are transfected with MHC genes which may be cloned from cells or tissue from the patient.
18 These transfected cells are then used to vaccinate the patient.

19 U.S. Patent No. 4,861,581 discloses a process comprising the steps of obtaining
20 monoclonal antibodies that are specific to an internal cellular component of neoplastic and
21 normal cells of the mammal but not to external components, labeling the monoclonal
22 antibody, contacting the labeled antibody with tissue of a mammal that has received
23 therapy to kill neoplastic cells, and determining the effectiveness of therapy by measuring

1 the binding of the labeled antibody to the internal cellular component of the degenerating
2 neoplastic cells. In preparing antibodies directed to human intracellular antigens, the
3 patentee recognizes that malignant cells represent a convenient source of such antigens.

4 U.S. Patent No. 5,171,665 provides a novel antibody and method for its production.
5 Specifically, the patent teaches formation of a monoclonal antibody which has the property
6 of binding strongly to a protein antigen associated with human tumors, e.g. those of the
7 colon and lung, while binding to normal cells to a much lesser degree.

8 U.S. Patent No. 5,484,596 provides a method of cancer therapy comprising
9 surgically removing tumor tissue from a human cancer patient, treating the tumor tissue to
10 obtain tumor cells, irradiating the tumor cells to be viable but non-tumorigenic, and using
11 these cells to prepare a vaccine for the patient capable of inhibiting recurrence of the
12 primary tumor while simultaneously inhibiting metastases. The patent teaches the
13 development of monoclonal antibodies which are reactive with surface antigens of tumor
14 cells. As set forth at col. 4, lines 45 et seq., the patentees utilize autochthonous tumor cells
15 in the development of monoclonal antibodies expressing active specific immunotherapy in
16 human neoplasia.

17 U.S. Patent No. 5,693,763 teaches a glycoprotein antigen characteristic of human
18 carcinomas and not dependent upon the epithelial tissue of origin.

19 U.S. Patent No. 5,783,186 is drawn to anti-Her2 antibodies which induce apoptosis
20 in Her2 expressing cells, hybridoma cell lines producing the antibodies, methods of
21 treating cancer using the antibodies and pharmaceutical compositions including said
22 antibodies.

1 U.S. Patent No. 5,849,876 describes new hybridoma cell lines for the production of
2 monoclonal antibodies to mucin antigens purified from tumor and non-tumor tissue
3 sources.

4 U.S. Patent No. 5,869,268 is drawn to a method for generating a human
5 lymphocyte producing an antibody specific to a desired antigen, a method for producing a
6 monoclonal antibody, as well as monoclonal antibodies produced by the method. The
7 patent is particularly drawn to the production of an anti-HD human monoclonal antibody
8 useful for the diagnosis and treatment of cancers.

9 U.S. Patent No. 5,869,045 relates to antibodies, antibody fragments, antibody
10 conjugates and single chain immunotoxins reactive with human carcinoma cells. The
11 mechanism by which these antibodies function is 2-fold, in that the molecules are reactive
12 with cell membrane antigens present on the surface of human carcinomas, and further in
13 that the antibodies have the ability to internalize within the carcinoma cells, subsequent to
14 binding, making them especially useful for forming antibody-drug and antibody-toxin
15 conjugates. In their unmodified form the antibodies also manifest cytotoxic properties at
16 specific concentrations.

17 U.S. Patent No. 5,780,033 discloses the use of autoantibodies for tumor therapy and
18 prophylaxis. However, this antibody is an anti-nuclear autoantibody from an aged
19 mammal. In this case, the autoantibody is said to be one type of natural antibody found in
20 the immune system. Because the autoantibody comes from "an aged mammal", there is no
21 requirement that the autoantibody actually comes from the patient being treated. In
22 addition the patent discloses natural and monoclonal antinuclear autoantibody from an

1 aged mammal, and a hybridoma cell line producing a monoclonal antinuclear
2 autoantibody.

3 U.S. Patent No. 5,916,561 discloses a specific antibody, VFF-18, and its variants
4 directed against the variant exon v6 of the CD44 gene. This antibody is an improvement
5 over the comparator antibody in that it recognizes a human CD44 v6 variant rather than a
6 rat CD44 v6 variant. In addition this antibody discloses diagnostic assays for CD44 v6
7 expression. There was no *in vitro* or *in vivo* function disclosed for this antibody.

8 U.S. Patent No. 5,616,468 discloses a monoclonal antibody, Var3.1, raised against
9 a synthetic peptide containing a sequence encoded by the human exon 6A of the CD44
10 gene. Specifically this antibody does not bind to the 90 kD form of human CD44 and is
11 distinguished from the Hermes-3 antibody. A method for detection of the v6 variant of
12 CD44 is provided, as well as a method for screening and assaying for malignant
13 transformation based on this antigen. A method for screening for inflammatory disease
14 based on detecting the antigen in serum is also provided.

15 U.S. Patent No. 5,879,898 discloses a specific antibody that binds to a 129 bp exon
16 of a human CD44 variant 6 that produces a 43 amino acid peptide. The monoclonal
17 antibody is produced by a number of hybridoma cell lines: MAK<CD44>M-1.1.12,
18 MAK<CD44>M-2.42.3, MAK<CD44>M-4.3.16. The antibody is generated from a fusion
19 protein that contains at least a hexapeptide of the novel CD44 v6 amino acid sequence.
20 Further, there is a disclosure of an immunoassay for the detection of exon 6 variant that
21 can be used as a cancer diagnostic. Significantly, there is no *in vitro* or *in vivo* function of
22 this antibody disclosed.

23 U.S. Patent No. 5,942,417 discloses a polynucleotide that encodes a CD44 like
24 polypeptide, and the method of making a recombinant protein using the polynucleotide and
McHale & Slavin, P.A.
2056.000025

1 its variants. Antibodies are claimed to these polypeptides however there are no specific
2 examples and there are no deposited clones secreting such antibodies. Northern blots
3 demonstrate the appearance of the polynucleotide in several types of tissues, but there is no
4 accompanying evidence that there is translation and expression of this polynucleotide.
5 Therefore, there is no evidence that there were antibodies to be made to the gene product
6 of this polynucleotide, that these antibodies would have either *in vitro* or *in vivo* function,
7 and whether they would be relevant to human cancerous disease.

8 U.S. Patent No. 5,885,575 discloses an antibody that reacts with a variant epitope
9 of CD44 and methods of identifying the variant through the use of the antibody. The
10 isolated polynucleotide encoding this variant was isolated from rat cells, and the antibody,
11 mAb1.1ASML, directed against this variant recognizes proteins of molecular weight 120
12 kD, 150 kD, 180 kD, and 200 kD. The administration of monoclonal antibody 1.1ASML
13 delayed the growth and metastases of rat BSp73ASML in isogenic rats. Significantly
14 1.1ASML does not recognize human tumors as demonstrated by its lack of reactivity to
15 LCLC97 human large-cell lung carcinoma. A human homolog was isolated from LCLC97
16 but no equivalent antibody recognizing this homolog was produced. Thus, although an
17 antibody specific to a variant of rat CD44 was produced and shown to affect the growth
18 and metastasis of rat tumors there is no evidence for the effect the this antibody against
19 human tumors. More specifically the inventors point out that this antibody does not
20 recognize human cancers.

21

22

23

1 Summary Of The Invention:

2 The instant inventors have previously been awarded U.S. Patent 6,180,357, entitled
3 "Individualized Patient Specific Anti-Cancer Antibodies" directed to a process for
4 selecting individually customized anti-cancer antibodies which are useful in treating a
5 cancerous disease. For the purpose of this document, the terms "antibody" and
6 "monoclonal antibody" (mAb) may be used interchangeably and refer to intact
7 immunoglobulins produced by hybridomas (e.g. murine or human), immunoconjugates
8 and, as appropriate, immunoglobulin fragments and recombinant proteins derived from
9 immunoglobulins, such as chimeric and humanized immunoglobulins, F(ab') and F(ab')₂
10 fragments, single-chain antibodies, recombinant immunoglobulin variable regions (Fv)s,
11 fusion proteins etc. It is well recognized in the art that some amino acid sequence can be
12 varied in a polypeptide without significant effect on the structure or function of the protein.
13 In the molecular rearrangement of antibodies, modifications in the nucleic or amino acid
14 sequence of the backbone region can generally be tolerated. These include, but are not
15 limited to, substitutions (preferred are conservative substitutions), deletions or additions.
16 Furthermore, it is within the purview of this invention to conjugate standard
17 chemotherapeutic modalities, e.g. radionuclides, with the CDMAB of the instant invention,
18 thereby focusing the use of said chemotherapeutics. The CDMAB can also be conjugated
19 to toxins, cytotoxic moieties, enzymes e.g. biotin conjugated enzymes, or hematogenous
20 cells.

21 This application utilizes substantially the method for producing patient specific
22 anti-cancer antibodies as taught in the '357 patent for isolating hybridoma cell lines which
23 encode for cancerous disease modifying monoclonal antibodies. These antibodies can be
24 made specifically for one tumor and thus make possible the customization of cancer

1 therapy. Within the context of this application, anti-cancer antibodies having either cell-
2 killing (cytotoxic) or cell-growth inhibiting (cytostatic) properties will hereafter be referred
3 to as cytotoxic. These antibodies can be used in aid of staging and diagnosis of a cancer,
4 and can be used to treat tumor metastases as well as primary tumors.

5 The prospect of individualized anti-cancer treatment will bring about a change in
6 the way a patient is managed. A likely clinical scenario is that a tumor sample is obtained
7 at the time of presentation, and banked. From this sample, the tumor can be typed from a
8 panel of pre-existing cancerous disease modifying antibodies. The patient will be
9 conventionally staged but the available antibodies can be of use in further staging the
10 patient. The patient can be treated immediately with the existing antibodies and/or a panel
11 of antibodies specific to the tumor can be produced either using the methods outlined
12 herein or through the use of phage display libraries in conjunction with the screening
13 methods herein disclosed. All the antibodies generated will be added to the library of anti-
14 cancer antibodies since there is a possibility that other tumors can bear some of the same
15 epitopes as the one that is being treated. The antibodies produced according to this method
16 may be useful to treat cancerous disease in any number of patients who have cancers that
17 bind to these antibodies.

18 Using substantially the process of US 6,180,357, the mouse monoclonal antibody
19 H460-16-2 was obtained following immunization of mice with cells from a patient's lung
20 tumor biopsy. The H460-16-2 antigen was expressed on the cell surface of a broad range
21 of human cell lines from different tissue origins. The breast cancer cell line MDA-MB-
22 231 (MB-231) and skin cancer cell A2058 were susceptible to the cytotoxic effects of
23 H460-16-2 *in vitro*.

1 The result of H460-16-2 cytotoxicity against MB-231 cells in culture was further
2 extended by its anti-tumor activity towards these cancer cells when transplanted into mice
3 (as disclosed in S.N. 10/603,000). Pre-clinical xenograft tumor models are considered
4 valid predictors of therapeutic efficacy.

5 In the preventative *in vivo* model of human breast cancer, H460-16-2 treatment was
6 significantly ($p<0.0001$) more effective in suppressing tumor growth during the treatment
7 period than an isotype control antibody, which was identical to H460-16-2 in structure and
8 size but incapable of binding MB-231 cells. At the end of the treatment phase, mice given
9 H460-16-2 had tumors that grew to only 1.3 percent of the control group. During the post
10 treatment follow-up period, the treatment effects of H460-16-2 were sustained and the
11 mean tumor volume in the treated groups continued to be significantly smaller than
12 controls until the end of the measurement phase. Using survival as a measure of antibody
13 efficacy, it was estimated that the risk of dying in the H460-16-2 treatment group was
14 about 71 percent of the antibody buffer control group ($p=0.028$) at 70 days post-treatment.
15 These data demonstrated that H40-16-2 treatment conferred a survival benefit compared to
16 the control-treated groups. H460-16-2 treatment appeared safe, as it did not induce any
17 signs of toxicity, including reduced body weight and clinical distress. Thus, H460-16-2
18 treatment was efficacious as it both delayed tumor growth and enhanced survival compared
19 to the control-treated groups in a well-established model of human breast cancer.

20 In addition H460-16-2 demonstrated anti-tumor activity against MB-231 cells in an
21 established *in vivo* tumor model (as outlined in S.N. 10/603,000). Treatment with H460-
22 16-2 was compared to the standard chemotherapeutic drug, cisplatin, and it was shown that
23 the cisplatin and H460-16-2 treatment groups had significantly ($p<0.001$) smaller mean
24 tumor volumes compared with groups treated with either antibody dilution buffer or the

1 isotype control antibody. H460-16-2 treatment mediated tumor suppression that was
2 approximately two-thirds that of cisplatin chemotherapy but without the significant
3 (19.2%) weight loss ($p<0.003$) and clinical distress, including 2 treatment-associated
4 deaths, observed with cisplatin treatment. The anti-tumor activity of H460-16-2 and its
5 minimal toxicity make it an attractive anti-cancer therapeutic agent.

6 In the post-treatment period, H460-16-2 showed a significant survival benefit
7 ($p<0.02$) as the risk of dying in the H460-16-2 group was about half of that in the isotype
8 control antibody group at >70 days after treatment. The observed survival benefit
9 continued past 120 days post-treatment where 100 percent of the isotype control and
10 cisplatin treated mice had died compared to 67 percent of the H460-16-2 treatment group.
11 H460-16-2 maintained tumor suppression by delaying tumor growth by 26 percent
12 compared to the isotype control antibody group. At 31 days post treatment, H460-16-2
13 limited tumor size by reducing tumor growth by 48 percent compared to the isotype
14 control group, which is comparable to the 49 percent reduction observed at the end of the
15 treatment. In the established tumor model of breast cancer, these results indicate the
16 potential of H460-16-2 to maintain tumor suppression beyond the treatment phase and
17 demonstrates the ability of the antibody to reduce the tumor burden and enhance survival
18 in a mammal.

19 In order to validate the H460-16-2 epitope as a drug target, the expression of
20 H460-16-2 antigen in normal human tissues was previously determined (S.N. 10/603,000).
21 This work was extended by comparison with an anti-CD44 antibody (clone L178). By
22 IHC staining with H460-16-2, the majority of the tissues again failed to express the H460-
23 16-2 antigen, including the cells of the vital organs, such as the liver, kidney (except for
24 marginal staining of tubular epithelial cells), heart, and lung. Results from tissue staining

1 indicated that H460-16-2 showed restricted binding to various cell types but had binding to
2 infiltrating macrophages, lymphocytes, and fibroblasts. The L178 antibody showed a
3 similar staining pattern. However, there were several differences of note; staining of
4 lymphocytes was more intense and had a wider distribution with L178 in comparison to
5 H460-16-2. Also, in one of the liver samples L178 stained the Kupffer cells while H460-
6 16-2 did not.

7 Localization of the H460-16-2 antigen and its prevalence within breast cancer
8 patients is important in assessing the benefits of H460-16-2 immunotherapy to patients and
9 designing effective clinical trials. To address H460-16-2 antigen expression in breast
10 tumors from cancer patients, tumor tissue samples from 50 individual breast cancer
11 patients were previously screened for expression of the H460-16-2 antigen (S.N.
12 10/603,000). Current work compared the staining of H460-16-2 to L178. The results of
13 the current study were similar to previous results and showed that 62 percent of tissue
14 samples stained positive for the H460-16-2 antigen while 76 percent of breast tumor
15 tissues were positive for the L178 epitope. Expression of H460-16-2 within patient
16 samples appeared specific for cancer cells as staining was restricted to malignant cells. In
17 contrast, H460-16-2 stained 4 of 10 samples of normal tissue from breast cancer patients
18 while L178 stained 6. Breast tumor expression of both the H460-16-2 and L178 antigen
19 appeared to be mainly localized to the cell membrane of malignant cells, making CD44 an
20 attractive target for therapy. H460-16-2 expression was further evaluated based on breast
21 tumor expression of the receptors for the hormones estrogen and progesterone, which play
22 an important role in the development, treatment, and prognosis of breast tumors. No
23 correlation was apparent between expression of the H460-16-2 antigen and expression of
24 the receptors for either estrogen or progesterone. When tumors were analyzed based on

1 their stage, or degree to which the cancer advanced, again there was no clear correlation
2 between H460-16-2 antigen expression and tumor stage. Similar results were obtained
3 with L178.

4 To further extend the potential therapeutic benefit of H460-16-2, the frequency and
5 localization of the antigen within various human cancer tissues was also previously
6 determined (S.N. 10/603,000). Current studies compared the staining of H460-16-2 to
7 clone L178. The majority of these tumor types were also positive for L178 antigen. As
8 with human breast tumor tissue, H460-16-2 and L178 localization occurred on the
9 membrane of tumor cells. However, there was substantially more membrane localization
10 with the L178 compared to the H460-16-2 antibody. Also, of the tumor types that were
11 stained by both H460-16-2 and L178, 43% of the tissues showed higher intensity staining
12 with the L178 antibody. There appears to be no form of CD44 that exactly matches the
13 IHC data presented herein based on comparisons with the IHC data from the literature.
14 The standard form of CD44 is normally expressed in the human brain; H460-16-2 antigen
15 is not. Antibodies directed against pan-CD44 isoforms do not stain the liver (including
16 Kupffer cells) and positively stain the endometrial glands in all phases of the reproductive
17 cycle. The H460-16-2 antigen is clearly present on Kupffer cells and is only present on
18 the secretory endometrial glands of the reproductive cycle. H460-16-2 antigen is clearly
19 present on tissue macrophages and only the variant forms V4/5 and V8/9 show occasional
20 macrophage staining. The similar yet distinct binding pattern seen with H460-16-2 in
21 comparison to anti-CD44 L178 indicates that the H460-16-2 antigen is an unique epitope
22 of CD44.

23 As outlined herein, additional biochemical data also indicate that the antigen
24 recognized by H460-16-2 is one of the forms of CD44. This is supported by studies

1 showing that a monoclonal antibody (L178) reactive against CD44 identifies proteins that
2 were bound to H460-16-2 by immunoprecipitation. Western blotting studies also suggest
3 that the epitope of CD44 recognized by H460-16-2 is not present on v6 or v10. The H460-
4 16-2 epitope is also distinguished by being carbohydrate and conformation dependent,
5 whereas many anti-CD44 antibodies are directed against peptide portions of CD44. These
6 IHC and biochemical results demonstrate that H460-16-2 binds to a variant of the CD44
7 antigen. Thus, the preponderance of evidence shows that H460-16-2 mediate anti-cancer
8 effects through ligation of an unique carbohydrate dependent conformational epitope
9 present on a variant of CD44.

10 *In toto*, this data demonstrates that the H460-16-2 antigen is a cancer associated
11 antigen and is expressed in humans, and is a pathologically relevant cancer target. Further,
12 this data also demonstrates the binding of the H460-16-2 antibody to human cancer tissues,
13 and can be used appropriately for assays that can be diagnostic, predictive of therapy, or
14 prognostic. In addition, the cell membrane localization of this antigen is indicative of the
15 cancer status of the cell due to the lack of expression of the antigen in most non-malignant
16 cells, and this observation permits the use of this antigen, its gene or derivatives, its protein
17 or its variants to be used for assays that can be diagnostic, predictive of therapy, or
18 prognostic.

19 Other studies, involving the use of anti-CD44 antibodies, have limitations of
20 therapeutic potential that is not exhibited by H460-16-2. H460-16-2 demonstrates both *in*
21 *vitro* and *in vivo* anti-tumor activity. Previously described antibodies such
22 MAK<CD44>M-1.1.12, MAK<CD44>M-2.42.3 and MAK<CD44>M-4.3.16 have no *in*
23 *vitro* or *in vivo* cytotoxicity ascribed to them and VFF-18 and Mab U36 shows no intrinsic

1 tumor cytotoxicity. In addition other anti-CD44 antibodies that have shown *in vivo* tumor
2 effects also have certain limitations that are not evident with H460-16-2. For example,
3 ASML1.1, K926, anti-CD44s and IM-78.1 show *in vivo* anti-tumor activity against rat,
4 murine, rat and murine tumors grown in xenograft models respectively. H460-16-2
5 demonstrates anti-tumor activity in a model of human cancer. H460-16-2 is also directed
6 against human CD44 while antibodies such as ASML1.1 recognize only rat CD44. The
7 clone 515 anti-CD44 antibody does inhibit peritoneal tumor implantation of a human
8 ovarian cell line but does not prevent or inhibit tumor growth. H460-16-2 is capable of
9 inhibiting human breast tumor growth in a SCID mouse xenograft model. GKW.A3 is an
10 anti-human CD44 monoclonal antibody capable of inhibiting tumor growth of a human
11 metastasizing melanoma grown in mice in a preventative but not an established model.
12 H460-16-2 has demonstrated significant anti-tumor activity in both preventative and
13 established murine xenograft models of human breast cancer. Consequently, it is quite
14 apparent that H460-16-2 has superior anti-tumor properties in comparison to previously
15 described anti-CD44 antibodies. It has demonstrated both *in vitro* and *in vivo* anti-tumor
16 activity on a human breast tumor in SCID mice and is directed against human CD44. It
17 also exhibits activity in a preventative and established (more clinically relevant) model of
18 human breast cancer.

19 In all, this invention teaches the use of the H460-16-2 antigen as a target for a
20 therapeutic agent, that when administered can reduce the tumor burden of a cancer
21 expressing the antigen in a mammal, and can also lead to a prolonged survival of the
22 treated mammal. This invention also teaches the use of a CDMAB (H460-16-2), and its
23 derivatives, to target its antigen to reduce the tumor burden of a cancer expressing the
24 antigen in a mammal, and to prolong the survival of a mammal bearing tumors that express

1 this antigen. Furthermore, this invention also teaches the use of detecting the H460-16-2
2 antigen in cancerous cells that can be useful for the diagnosis, prediction of therapy, and
3 prognosis of mammals bearing tumors that express this antigen.

4 If a patient is refractory to the initial course of therapy or metastases develop, the
5 process of generating specific antibodies to the tumor can be repeated for re-treatment.
6 Furthermore, the anti-cancer antibodies can be conjugated to red blood cells obtained from
7 that patient and re-infused for treatment of metastases. There have been few effective
8 treatments for metastatic cancer and metastases usually portend a poor outcome resulting
9 in death. However, metastatic cancers are usually well vascularized and the delivery of
10 anti-cancer antibodies by red blood cells can have the effect of concentrating the antibodies
11 at the site of the tumor. Even prior to metastases, most cancer cells are dependent on the
12 host's blood supply for their survival and anti-cancer antibody conjugated to red blood
13 cells can be effective against *in situ* tumors as well. Alternatively, the antibodies may be
14 conjugated to other hematogenous cells, e.g. lymphocytes, macrophages, monocytes,
15 natural killer cells, etc.

16 There are five classes of antibodies and each is associated with a function that is
17 conferred by its heavy chain. It is generally thought that cancer cell killing by naked
18 antibodies are mediated either through antibody-dependent cell-mediated cytotoxicity
19 (ADCC) or complement-dependent cytotoxicity (CDC). For example murine IgM and
20 IgG2a antibodies can activate human complement by binding the C-1 component of the
21 complement system thereby activating the classical pathway of complement activation
22 which can lead to tumor lysis. For human antibodies, the most effective complement
23 activating antibodies are generally IgM and IgG1. Murine antibodies of the IgG2a and

1 IgG3 isotype are effective at recruiting cytotoxic cells that have Fc receptors which will
2 lead to cell killing by monocytes, macrophages, granulocytes and certain lymphocytes.
3 Human antibodies of both the IgG1 and IgG3 isotype mediate ADCC.

4 Another possible mechanism of antibody mediated cancer killing may be through
5 the use of antibodies that function to catalyze the hydrolysis of various chemical bonds in
6 the cell membrane and its associated glycoproteins or glycolipids, so-called catalytic
7 antibodies.

8 There are two additional mechanisms of antibody mediated cancer cell killing
9 which are more widely accepted. The first is the use of antibodies as a vaccine to induce
10 the body to produce an immune response against the putative antigen that resides on the
11 cancer cell. The second is the use of antibodies to target growth receptors and interfere
12 with their function or to down regulate that receptor so that effectively its function is lost.

13 Accordingly, it is an objective of the invention to utilize a method for producing
14 cancerous disease modifying antibodies from cells derived from a particular individual
15 which are cytotoxic with respect to cancer cells while simultaneously being relatively non-
16 toxic to non-cancerous cells, in order to isolate hybridoma cell lines and the corresponding
17 isolated monoclonal antibodies and antigen binding fragments thereof for which said
18 hybridoma cell lines are encoded.

19 It is an additional objective of the invention to teach methods of utilizing the
20 isolated monoclonal antibody or antigen binding fragment thereof encoded by the clone
21 deposited with the ATCC as PTA-4621 for determining a presence of cells which express a
22 CD44 antigenic moiety which specifically binds to an isolated monoclonal antibody or

1 antigen binding fragment thereof encoded by the clone deposited with the ATCC as PTA-
2 4621.

3 It is yet a further objective of the instant invention to teach methods for enhancing
4 the survival of a patient having a cancerous disease via the use of an isolated monoclonal
5 antibody or antigen binding fragment thereof encoded by the clone deposited with the
6 ATCC as PTA-4621, which antibody specifically binds to a CD44 antigenic moiety.

7 It is an additional objective of the invention to teach CDMAB and antigen binding
8 fragments thereof.

9 It is a further objective of the instant invention to produce CDMAB whose
10 cytotoxicity is mediated through ADCC.

11 It is yet an additional objective of the instant invention to produce CDMAB whose
12 cytotoxicity is mediated through CDC.

13 It is still a further objective of the instant invention to produce CDMAB whose
14 cytotoxicity is a function of their ability to catalyze hydrolysis of cellular chemical bonds.

15 A still further objective of the instant invention is to produce CDMAB which are
16 useful in a binding assay for diagnosis, prognosis, and monitoring of cancer.

17 Other objects and advantages of this invention will become apparent from the
18 following description wherein are set forth, by way of illustration and example, certain
19 embodiments of this invention.

20

21 Brief Description of the Figures:

1 The patent or application file contains at least one drawing executed in color. Copies of
2 this patent or patent application publication with color drawing(s) will be provided by the
3 Office upon request and payment of the necessary fee.

4 Figure 1. Western blot of MDA-MB-468 membranes probed with H460-16-2. Lane 1:
5 Membrane proteins separated under reducing conditions. Lane 2: Membrane proteins
6 separated under non-reducing conditions. Molecular weight markers are indicated on the
7 left.

8 Figure 2. Western blot of membranes probed with H460-16-2. Lane 1: MDA-MB-468
9 membranes. Lane 2: MDA-MB-231 membranes. Molecular weight markers are indicated
10 on the left.

11 Figure 3. 2-Dimensional Western blot and SDS-PAGE of MDA-MB-468 membrane
12 proteins. Panel A demonstrates the position of the 2 proteins recognized by H460-16-2.
13 Panel B demonstrates a similar blot probed with an isotype control antibody. Panel C
14 shows a SYPRO Ruby-stained gel of MDA-MB-468 membranes. Arrows indicate the
15 position of protein spots corresponding to panel A.

16 Figure 4. 2-Dimensional Western blot of MDA-MB-231 membrane proteins. The major
17 binding protein is indicated with an arrow.

18 Figure 5. Effect of deglycosylation on the binding of H460-16-2 to MDA-MB-231
19 membranes. Panel A demonstrates the binding of H460-16-2 in a Western blot to
20 untreated MDA-MB-231 cell membranes (Lane 2), membranes treated with glycosidases
21 (see text) at 37° C for 24 hr (Lane 3), and membranes treated with glycosidases at 25° C for
22 24 hr (Lane 4). Lane 1 shows the position of the molecular weight markers. Panel B
23 demonstrates the binding of the high-mannose binding lectin GNA to a similar blot.

1 Figure 6. SDS-PAGE (Panel A) and Western blot (Panel B) of MDA-MB-231 membrane
2 proteins immunoprecipitated with H460-16-2. Lane 1: Total MDA-MB-231 membrane
3 proteins Lane 2: H460-16-2 immunoprecipitated proteins. Arrow demonstrates 80-90 kD
4 H460-16-2 binding protein.

5 Figure 7. Western blots of proteins probed with H460-16-2 (Panel A), anti-CD44 (clone
6 L178, Panel B) and anti-HSP90 (Panel C). Lane 1: Total MDA-MB-231 membrane
7 proteins; Lane 2: H460-16-2 immunoprecipitated proteins; Lane 3: Molecular weight
8 standards.

9 Figure 8. Western blots of proteins probed with H460-16-2 (Panel A, Lane 2), anti-CD44
10 (clone L178, Panel B, Lane 2), anti-CD44var6 (clone VFF-7, Panel C, Lane 2) and anti-
11 CD44var10 (clone VFF-14, Panel D, Lane 2). Lane 1 of each blot contains the molecular
12 weight standards.

13

14 Figure 9. Representative histograms for H460-16-2 binding by FACS. Histograms for
15 binding of H460-16-2 (20 μ g/mL), 107.3 isotype control (20 μ g /mL) and anti-EGFR (5 μ g
16 /mL) are presented for breast cancer (MDA-MB-231 and MDA-MB-468) and normal
17 (Hs578.Bst and CCD-27sk) cell lines.

18 Figure 10. Representative micrographs showing the binding pattern obtained with H460-
19 16-2 (A) and the anti-CD44 (L178) antibody (B) on tissues sections of tonsil from a
20 normal human tissue array. There is more intense and widely distributed staining of
21 lymphocytes with L178 than with ARH460-16-2 where the staining is more limited to
22 mantle zone of lymphoid nodules (black arrow), leaving the germinal center (green arrow)
23 with weak staining. Magnification is 200X.

1 Figure 11. Representative micrographs showing the binding pattern obtained with H460-
2 16-2 (A) and the anti-CD44 (L178) antibody (B) on tissues sections of liver from a normal
3 human tissue array. There is staining for Kupffer cells (arrows) in hepatic sinusoids with
4 L178 and not with H460-16-2. Magnification is 200X.

5 Figure 12. Representative micrograph of H460-16-2 binding to breast cancer tumor
6 (infiltrating duct carcinoma). The yellow and orange arrows in panel point to stromal cells
7 and sheets of malignant cells respectively. Magnification is 100X.

8 Figure 13. Representative micrographs showing the binding pattern obtained with H460-
9 16-2 (A) and the anti-CD44 (L178) antibody (B) on paget's disease breast tissue sections
10 from a human breast cancer tissue array. There is a membranous staining of malignant
11 cells (arrows) with L178 versus negative staining with H460-16-2. Magnification is 200X.

12 Figure 14. Representative micrographs showing the binding pattern obtained with H460-
13 16-2 (A) and the anti-CD44 (L178) antibody (B) on uterine cervix squamous cell
14 carcinoma tissue sections from a human multi-tumor tissue array. There is a stronger
15 membranous staining of malignant cells with H460-16-2 than with L178. Magnification is
16 200X.

17 Figure 15. Representative micrographs showing the binding pattern obtained with H460-
18 16-2 (A) and the anti-CD44 (L178) antibody (B) on adenocarcinoma lung tissue sections
19 from a human multi-tumor tissue array. There is +++ scoring of malignant cells (arrows)
20 with L178 versus +/- with H460-16-2. Magnification is 200X.

21

22 Detailed Description Of The Invention:

23 Example 1

1 Identification of Binding Proteins by Western Blotting

2 To identify the antigen(s) recognized by the antibody H460-16-2, cell membranes
3 expressing this antigen were subjected to gel electrophoresis, and transferred to
4 membranes. Western blotting was used to determine proteins detected by this antibody.

5

6 1. Membrane Preparation

7 Previous work demonstrated binding by FACS of H460-16-2 to the breast cancer
8 cell lines MDA-MB-231 (MB-231) and MDA-MB-468 (MB-468). Accordingly,
9 membrane preparations from these two cell lines were used for antigen identification.
10 Total cell membranes were prepared from confluent cultures of MB-231 or MB-468 breast
11 cancer cells. Media was removed from flasks, and the cells were washed 3 times with
12 PBS. After the final wash, cells were dissociated with Dissociation Buffer (Gibco-BRL;
13 Grand Island, NY) for 5 min at 37°C. Cells were collected and centrifuged at 1200 rpm for
14 10 minutes at 4°C. After centrifugation, cell pellets were resuspended in 1 mL of
15 hypotonic lysis buffer containing 10 µg/mL leupeptin, 10 µg/mL aprotinin and 25 µg/mL
16 4-(2-aminoethyl)-benzenesulfonyl fluoride. Cells were then lysed using 5 cycles of rapid
17 freezing and thawing. Cell lysates were centrifuged at 9500 rpm for 10 min at 4°C to
18 remove nuclear particulate. Supernatant was harvested, and then centrifuged at 75,000 x g
19 for 57 min, at 4°C. Supernatant was carefully removed from tubes, and pellets were
20 resuspended in 0.5 to 1 mL of hypotonic lysis buffer containing 1 percent Triton X-100.
21 Membranes were then assayed for protein content, and stored at -80°C.

22

23 2. 1-Dimensional SDS-PAGE

1 Membrane proteins were separated by 1-dimensional SDS-polyacrylamide gel
2 electrophoresis. 20 µg of membrane protein was loaded onto a lane of a 12 percent SDS-
3 PAGE gel. A sample of pre-stained molecular weight markers (Biorad; Mississauga, ON)
4 was run in a reference lane. The sample was separated by electrophoresis under non-
5 reducing conditions, in the absence of dithiothreitol (DTT). Electrophoresis was carried
6 out at 100 V for 10 min, followed by 65 min at 150 V. Proteins were transferred from the
7 gel to PVDF (Millipore; Billerica, MA) membranes by electroblotting for 16 hr at 40 V.
8 Quantitative transfer was assessed by noting complete transfer of the prestained markers
9 from the gel to the membrane.

10 After transfer, membranes were blocked with 5 percent skim milk powder in Tris-
11 buffered saline containing 0.5 percent Tween (TBST) for 2 hr. Membranes were then
12 incubated with 2 – 2.5 µg/mL H460-16-2 diluted into 3 percent skim milk powder in TBST
13 for 2 hr. After washing 3 times with TBST, membranes were incubated with goat anti-
14 mouse IgG (Fc) conjugated to horseradish peroxidase (HRP) from Jackson
15 Immunologicals (West Grove, PA) for 1 hr. This incubation was followed by washing 3
16 times with TBST, followed by incubation with the HRP substrate TMB (substrate kit from
17 Vector Laboratories; Burlington, ON).

18 Figure 1 demonstrates the results of the Western blotting as described. H460-16-2
19 binds clearly to 2 molecular weight (MW) regions of the separated MB-468 membrane
20 proteins (Lane 2). By comparison to the molecular weight standards, the antibody binds to
21 proteins of MW 80-90 kD and MW 120-150 kD. The epitope recognized by the antibody
22 H460-16-2 appears to be to a conformational epitope, since the antibody was unable to
23 bind spots transferred from gels under reducing conditions in the presence of DTT (Lane
24 1). Figure 2 compares the binding of H460-16-2 to membranes from MB-468 (Lane 1)

1 and MB-231 (Lane 2) cells. In MB-468 membranes, H460-16-2 binds to both the 80-90
2 kD and 120-150 kD proteins with equal intensity. In MB-231 membranes, the major
3 binding protein is the 80-90 kD protein, with a less intense binding protein identified in the
4 120-150 kD range.

5 6 **3. 2-Dimensional SDS-PAGE**

7 In order to obtain better resolution of the binding entities, and to further
8 characterize the proteins, 2-dimensional electrophoresis was carried out. Total membrane
9 proteins (75-200 µg) prepared as described above, were precipitated using the PlusOne 2-D
10 Clean Up kit (Amersham; Baie D'Urfé, QC), and then resuspended in rehydration buffer
11 containing ampholytes in the pH range 3-10. Samples were centrifuged to remove
12 particulate material, and then loaded onto IPG strips (Amersham; Baie D'Urfé, QC) in the
13 presence of a rehydration solution. Proteins were focussed using the following protocol:
14 16 hr for rehydration; 500 V, 250 Vhrs, 1000 V, 500 Vhrs; 5000 V, 7500 V hrs. The strip
15 was then removed from the strip holders, and incubated in an SDS-PAGE equilibration
16 buffer. After 15 min, the strip was placed on the top of an 8 percent gel, and sealed with
17 an agarose solution. Prestained MW markers were loaded beside the strip. Electrophoresis
18 was carried out at 100 V for 10 min, followed by 65 min at 150 V. One of the gels was
19 fixed for 30 min with 10 percent methanol/7 percent acetic acid, and then stained with the
20 fluorescent dye SYPRO RubyTM (Molecular Probes, Eugene, OR). Protein spots were
21 visualized under UV light. From a second and third gel, proteins were transferred from the
22 gels to PVDF (Millipore) membranes by electroblotting for 16 hr at 40 V. Quantitative
23 transfer was assessed by determining the complete transfer of the prestained markers from
24 the gel to the membrane.

1 After transfer, membranes were blocked with 5 percent skim milk powder in Tris-
2 buffered saline containing 0.5 percent Tween (TBST) for 2 hr. One of the membranes was
3 then incubated with 2–2.5 µg/mL H460-16-2 diluted into 3 percent skim milk powder in
4 TBST for 2 hrs. A similar membrane was incubated with the same concentration of an
5 isotype control (Mouse anti-trinitrophenol, IgG1,κ; clone 107.3 (BD Biosciences;
6 Oakville, ON). After washing 3 times with TBST, membranes were incubated with goat
7 anti-mouse IgG (Fc) conjugated to horseradish peroxidase (HRP) from Jackson
8 Immunologicals (West Grove, PA) for 1 hr. This incubation was followed by washing 3
9 times with TBST, followed by incubation with the HRP substrate TMB (substrate kit from
10 Vector Laboratories; Burlington, ON).

11 Figure 3a demonstrates the Western blot obtained from MB-468 membranes
12 incubated with H460-16-2. Two distinct binding spots are observed, with molecular
13 weights corresponding with those obtained by 1-dimensional electrophoresis. One is
14 observed at a MW of approximately 80-90 kD according to the MW standards, and is in
15 the acidic portion of the gel with an estimated pI of 3 – 4. The second spot is in the MW
16 range of 120-150 kD according to the MW standards, and has a pI more basic than the 80-
17 90 kD protein. Figure 3b demonstrates the Western blot obtained from membranes
18 incubated with the isotype control antibody. No spots were visible on this blot, indicating
19 that the binding of H460-16-2 was not due to non-specific binding. Figure 3c shows a
20 SYPRO RubyTM stained 2D-gel of MB-468 membrane proteins. Note that when a similar
21 Western blot is run with MB-231 membranes, only the 80-90 kD spot is observed (Figure
22 4).

23 Example 2

Determining Glycosylation of Antigens Bound by H460-16-2

In order to determine if the antigen(s) recognized by the antibody H460-16-2 were glycoproteins, MB-231 membranes were incubated with PNGase F, Endo-o-glycosidase, and sialidase A according to manufacturer's protocol (DeglycoPro deglycosylation kit; Prozyme, San Leandro, CA) for 24 hr at room temperature or at 37 °C. Membranes were separated by 1-D polyacrylamide gel electrophoresis as described, and then Western blotting was carried out as described with H460-16-2. The results of the Western blot are shown in Figure 5. In MB-468 membranes that were not treated with glycosidases, H460-16-2 recognized the expected 85-95 kD band (Figure 5, Panel A, Lane 2). In membranes treated with glycosidases at 25 °C, there is a distinct shift of this band to a lower molecular weight (Lane 4). In membranes treated with glycosidases at 37 °C, the binding of H460-16-2 is eliminated (Lane 3). In order to determine the completeness of deglycosylation, a similar blot was probed with the high-mannose binding lectin galanthus nivalis agglutinin (GNA). Results observed in Figure 5, Panel B demonstrate that deglycosylation is incomplete at 25°C (Lane 4) and essentially complete at 37°C (Lane 3). Thus, under conditions of complete deglycosylation, H460-16-2 is unable to bind to its antigen.

Taken together, these results suggest that the 80-90 kD band is a glycoprotein. In addition, these results present evidence that the epitope recognized by H460-16-2 is carbohydrate-dependent.

Example 3

1 Identification of Antigens Bound by H460-16-2

2 1. Immunoprecipitation

3
4 1 mL of Protein G Dynabeads (DYNAL) was washed 3 times with 0.1 M sodium
5 phosphate buffer, pH 6.0. 2500 µg of H460-16-2 was added to the washed beads in a total
6 volume of 500 µl. The mixture was incubated with gentle mixing for 1 hr. Unbound
7 antibody was removed and the H460-16-2-coated beads were washed 3 times in 2.5 mL
8 volumes of 0.1 M sodium phosphate buffer, pH 7.4 containing 0.1 percent Tween-20. The
9 H460-16-2-coated beads were washed 2 times in 5 mL of 0.2 M triethanolamine, pH 8.2
10 followed by the addition of 5 mL. H460-16-2 was chemically crosslinked to the beads by
11 gentle mixing in the presence of 5 mL of 0.2 M triethanolamine, pH 8.2 containing 20 mM
12 dimethyl pimelimidate for 30 min. The reaction was stopped by the addition of 5 mL of 50
13 mM Tris, pH 7.5. After 15 min incubation, the H460-16-2 crosslinked beads were washed
14 3 times in PBS containing 0.1 percent Tween-20. The H460-16-2-crosslinked beads were
15 pre-eluted by incubation with 0.1 M citrate pH 3 for 3 min followed by 3 washes in 0.1 M
16 sodium phosphate buffer, pH 7.4 containing 0.1 percent Tween-20.

17 Three mg of total membrane protein from MB-231 cells was incubated with H460-
18 16-2 chemically crosslinked beads in 0.1 M sodium phosphate buffer, pH 7.4 containing
19 0.1 percent Tween-20, 5 percent glucose, 5 percent mannose, 5 percent galactose and
20 protease inhibitors at 4°C for 4 hr. After incubation, the immunoprecipate was washed 3
21 times in PBS containing 150 mM NaCl and 0.1 percent Tween-20. Protein was eluted
22 from the beads by incubating the H460-16-2-crosslinked beads with 0.1 M citrate, pH 3 for
23 4 min. Eluted protein was stored at -80°C. Immunoprecipitated protein from 3 mg of
24 protein was loaded onto a single lane of an 8 percent non-reducing SDS-PAGE gel. A

sample of pre-stained molecular weight markers (Biorad; Mississauga, ON) was run in a reference lane. The sample was separated by electrophoresis at 100 V for 10 min, followed by 65 min at 150 V. Proteins were stained with SYPRO Ruby™.

In a parallel Western blot, proteins immunoprecipitated as described from 100 µg of MB-231 membrane proteins were separated by electrophoresis. Proteins were transferred from the gel to PVDF (Millipore; Billerica, MA) membranes by electroblotting for 16 hr at 40 V. Quantitative transfer was assessed by noting complete transfer of the prestained markers from the gel to the membrane.

After transfer, the membrane was blocked with 5 percent skim milk powder in Tris-buffered saline containing 0.5 percent Tween (TBST) for 2 hr. The membrane was probed with 5 µg/ml H460-16-2 diluted into 3 percent skim milk powder in TBST for 2 hr. After washing 3 times with TBST, membranes were incubated with appropriate secondary antibody: goat anti-mouse IgG (Fc) conjugated to horseradish peroxidase (HRP) from Jackson Immunologicals (West Grove, PA) for 1 hr. This incubation was followed by washing 3 times with TBST, followed by incubation with the HRP substrate TMB (substrate kit from Vector Laboratories; Burlington, ON).

Figure 6 depicts the gel and blot obtained from the proteins immunoprecipitated by H460-16-2. On the gel (Panel A) there is a band in the 80-90 kD region in the lane containing the immunoprecipitated proteins (see arrow, Lane 2). The high molecular weight bands are comprised of intact antibodies. No other proteins are present in the sample. In a corresponding Western blot (Panel B), H460-16-2 reacts strongly with the immunoprecipitated proteins (Lane 2), with a similar binding profile to that seen in the total membranes (Lane 1).

2. Mass Spectroscopy

The region of the gel corresponding to the 80-90 kD protein immunoprecipitated by H460-16-2 (Figure 6, Panel A, Lane 2) was excised using a sterile pipette tip. This gel plug was then used for identification of proteins by mass spectroscopy.

The sample was subjected to robotic in-gel digestion using trypsin (ProGest) and a portion of the resulting digest supernatant was used for MALDI/MS analysis. Spotting was performed robotically (ProMS) with ZipTips; peptides were eluted from the C18 material with matrix (α -cyano 4-hydroxy cinnamic acid) prepared in 60 percent acetonitrile, 0.2 percent TFA. The sample was analyzed by nano LC/MS/MS on a Micromass Q-Tof2 using a 75 μ m C18 column at a flow-rate of 200 nL/min. The MS/MS data were searched using a local copy of MASCOT.

The proteins identified by LC/MS/MS analysis of the H460-16-2 immunoprecipitated material are presented in Table 1. A score was assigned which is a composite score based on the number of peptides matched, and the level of identity.

Table 1. Proteins Identified by H460-16-2 Immunoprecipitation of MDA-MB-231 Membranes

Observed MW	Method	Protein ID	Score	NCBI Accession #
80-90 KD	LC/MS/MS	CD44	239	gi 2134882

The only protein present in the sample was matched to human CD44, an 80 kD glycoprotein present on the cell surface of lymphocytes and multiple types of cancer cells.

3. Confirmation

1 Confirmation of the putative antigen was carried out by determining whether a
2 known anti-CD44 monoclonal antibody would react with the protein immunoprecipitated
3 by H460-16-2. Total MB-231 membrane proteins and H460-16-2-immunoprecipitated
4 proteins were separated by 1-dimensional SDS-PAGE. Electrophoresis and Western
5 blotting were carried out as described above. Membranes were incubated with 5-10 µg/ml
6 H460-16-2, anti-CD44 (clone L178, BD Biosciences, Oakville, ON) or anti-HSP90
7 (negative control; clone 16F1 (Stressgen, Victoria, BC)) diluted into 3 percent skim milk
8 powder in TBST for 2 hr. Figure 7 demonstrates the results of the Western blotting as
9 described. Panel A shows binding of H460-16-2 to total membrane proteins (Lane 1) and
10 immunoprecipitated proteins (Lane 2). The major binding protein is in the 80-90 kD
11 region, although a band in the 120-150 kD region can also be seen in the total membrane
12 proteins. When a parallel blot was probed with L178 (Panel B), a similar pattern was seen.
13 L178 bound strongly to the protein immunoprecipitated by H460-16-2. The binding
14 pattern of H460-16-2 and L178 to the total membrane proteins is very similar (Panel A and
15 B, Lane 1). A parallel blot probed with the negative control anti-HSP90 (Panel C) showed
16 that this antibody did not bind to the immunoprecipitated material, although it did bind to a
17 distinct 90 kD protein in the total membrane preparation. This result confirms that the
18 binding of anti-CD44 to the immunoprecipitated protein was specific, and not due to the
19 presence of contaminating proteins since HSP90 is an ubiquitous and “sticky” cellular
20 chaperone protein.

21 Figure 8 presents the results of an experiment carried out to compare the specificity
22 of H460-16-2 to known anti-CD44 antibodies. MDA-MB-231 membranes were separated
23 by electrophoresis and transferred to PVDF membranes as described. Membranes were
24 probed with H460-16-2 (Panel A), anti-CD44 (clone L178, BD Biosciences; Panel B),

1 anti-CD44var6 (clone VFF-7, Bender Medsystems (San Bruno, CA); Panel C) and anti-
2 CD44var10 (clone VFF-14, Bender Medsystems; Panel D). H460-16-2 and L178 react
3 with an 80-90 kD and 120-150 kD protein, and have identical binding patterns by Western
4 blot. By contrast, antibodies specific for CD44 variants 6 and 10 demonstrate differential
5 binding from H460-16-2 and each other. Both variant antibodies bind to a wider range of
6 proteins, and do not show strong binding to the 80 kD region of the gel. It is therefore
7 unlikely that H460-16-2 is directed towards either variant 6 or variant 10.

8 The mass spectroscopic identification, as well as the confirmation using a known
9 antibody, demonstrates that the antigen for H460-16-2 is a form of CD44. This is also
10 consistent with the deglycosylation experiments in Example 2, as CD44 may contain
11 around 50 percent N-linked sugars by weight. These experiments also indicate that H460-
12 16-2 binds to a carbohydrate-dependent epitope of CD44.

13

14 Example 4

15 The hybridoma cell line H460-16-2 was deposited, in accordance with the
16 Budapest Treaty, with the American Type Culture Collection, 10801 University Blvd.,
17 Manassas, VA 20110-2209 on September 4, 2002, under Accession Number PTA-4621.
18 In accordance with 37 CFR 1.808, the depositors assure that all restrictions imposed on the
19 availability to the public of the deposited materials will be irrevocably removed upon the
20 granting of a patent.

21 **Antibody Production:**

22 H460-16-2 monoclonal antibody was produced by culturing the hybridomas in CL-
23 1000 flasks (BD Biosciences, Oakville, ON) with collections and reseeding occurring

twice/week and standard antibody purification procedures with Protein G Sepharose 4 Fast Flow (Amersham Biosciences, Baie d'Urfé, QC). It is within the scope of this invention to utilize monoclonal antibodies that are human, humanized, chimerized or murine antibodies. H460-16-2 was compared to a number of both positive (anti-Fas (EOS9.1, IgM, kappa, 10 µg/mL, eBioscience; San Diego, CA), anti-Her2/neu (IgG1, kappa, 10 µg/mL, Inter Medico; Markham, ON), anti-EGFR (C225, IgG1, kappa, 5 µg/mL, Cedarlane; Hornby, ON), Cycloheximide (100 µM, Sigma; Oakville, ON), NaN₃ (0.1 percent, Sigma; Oakville, ON)) and negative (107.3 (anti-TNP, IgG1, kappa, 20 µg/mL, BD Biosciences; Oakville, ON), G155-178 (anti-TNP, IgG2a, kappa, 20 µg/mL, BD Biosciences; Oakville, ON), MPC-11 (antigenic specificity unknown, IgG2b, kappa, 20 µg/mL, BD Biosciences; Oakville, ON), J606 (anti-fructosan, IgG3, kappa, 20 µg/mL, BD Biosciences; Oakville, ON), IgG Buffer (2 percent)) controls in a cytotoxicity assay (Table 3). Breast cancer (MB-231, MB-468), melanoma (A2058, A375), colon cancer (HT-29), lung cancer (NCI-H460, A549), ovarian cancer (OVCAR-3), prostate cancer (PC-3), and non-cancer (CCD-27sk, Hs578.Bst, Hs888.Lu) cell lines were tested (all from the ATCC, Manassas, VA). The Live/Dead cytotoxicity assay was obtained from Molecular Probes (Eugene, OR). The assays were performed according to the manufacturer's instructions with the changes outlined below. Cells were plated before the assay at the predetermined appropriate density. After 2 days, 100 µL of purified antibody was diluted into media, and then were transferred to the cell plates and incubated in a 5 percent CO₂ incubator for 5 days. The plate was then emptied by inverting and blotted dry. Room temperature DPBS containing MgCl₂ and CaCl₂ was dispensed into each well from a multichannel squeeze bottle, tapped 3 times, emptied by inversion and then blotted dry. 50 µL of the fluorescent Live/Dead dye

1 diluted in DPBS containing $MgCl_2$ and $CaCl_2$ was added to each well and incubated at
2 $37^\circ C$. in a 5 percent CO_2 incubator for 30 min. The plates were read in a Perkin-Elmer
3 HTS7000 fluorescence plate reader and the data was analyzed in Microsoft Excel and the
4 results were tabulated in Table 2. The data represented an average of 4 experiments tested
5 in triplicate and presented qualitatively in the following fashion: 4/4 experiments greater
6 than threshold cytotoxicity (+++), 3/4 experiments greater than threshold cytotoxicity (++)
7 2/4 experiments greater than threshold cytotoxicity (+). Unmarked cells in Table 2
8 represented inconsistent or effects less than the threshold cytotoxicity. The H460-16-2
9 antibody produced selective cytotoxicity in A2058 melanoma cells and MB-231 breast
10 cancer cells but did not produce cytotoxicity against the remaining cancer cells,
11 demonstrating properties of specific cytotoxicity towards cancer cells. Importantly the
12 isolated antibody did not produce cytotoxicity against a number of non-cancer cells such as
13 CCD-27sk, Hs578.Bst or Hs888.Lu. The chemical cytotoxic agents induced their expected
14 cytotoxicity while a number of other antibodies which were included for comparison also
15 performed as expected given the limitations of biological cell assays.

Table 2: *In Vitro* Cytotoxicity

	Melanoma		Breast		Lung		Colon	Ovary	Prostate	Normal		
	A2058	A375	MB-231	MB-468	NCI-H460	A549	HT-29	OVCAR-3	PC-3	CCD-27sk	t	Hs888.Lu
H460-16-2 (20 $\mu g/mL$)	+		+									
Positive Controls												
Anti-Fas (20 $\mu g/mL$)	++	+++						+++				
Anti-Her2/neu (10 $\mu g/mL$)				+				++				
Anti-EGFR (c528, 5 $\mu g/mL$)												
Cycloheximide (100 μM)	+++	+++	+++	+++	++	+++	+++	+++	+++	+++	+	+++
Negative Controls												
IgG1 (107.3, 20 $\mu g/mL$)									+			
Human IgG (10 $\mu g/mL$)												
IgG Buffer (2%)												

16

17 Cells were prepared for FACS by initially washing the cell monolayer with DPBS
18 (without Ca^{++} and Mg^{++}). Cell dissociation buffer (INVITROGEN; Burlington, ON) was

1 then used to dislodge the cells from their cell culture plates at 37°C. After centrifugation
2 and collection the cells were resuspended in DPBS containing MgCl₂, CaCl₂ and 25 percent
3 fetal bovine serum at 4°C (wash media) and counted, aliquoted to appropriate cell density,
4 spun down to pellet the cells and resuspended in staining media (DPBS containing MgCl₂,
5 CaCl₂ and 2 percent fetal bovine serum) at 4°C in the presence of test antibodies (H460-16-
6 2) or control antibodies (isotype control, anti-Her2/neu or anti-EGFR) at 20 µg/mL on ice
7 for 30 minutes. Prior to the addition of Alexa Fluor 488-conjugated secondary antibody the
8 cells were washed once with wash media. The Alexa Fluor 488-conjugated antibody in
9 staining media was then added for 20 min. The cells were then washed for the final time
10 and resuspended in staining media containing 1 µg /mL propidium iodide. Flow cytometric
11 acquisition of the cells was assessed by running samples on a FACScan using the
12 CellQuest software (BD Biosciences; Oakville, ON). The forward (FSC) and side scatter
13 (SSC) of the cells were set by adjusting the voltage and amplitude gains on the FSC and
14 SSC detectors. The detectors for the three fluorescence channels (FL1, FL2, and FL3) were
15 adjusted by running cells stained with purified isotype control antibody followed by Alexa
16 Fluor 488-conjugated secondary antibody such that cells had a uniform speak with a
17 median fluorescent intensity of approximately 1-5 units. Live cells were acquired by gating
18 for FSC and propidium iodide exclusion. For each sample, approximately 10,000 live cells
19 were acquired for analysis and the results presented in Table 3.

20 Table 3 tabulated the mean fluorescence intensity fold increase above isotype
21 control and is presented qualitatively as: less than 5 (-); 5 to 50 (+); 50 to 100 (++); above

Table 3: FACS Analysis

Antibody	NCI-H460	A549	Hs888.Lu	HT-29	PC-3	OVCAR-3	A375	A2058	CCD-27sk	MB-231	MB-468	Hs578.Bst
Anti-EGFR	+(100%)	+(100%)	++	++	+(98%)	++	+(99%)	-	+(98%)	+++	+++	+(97%)
Anti-HER2/neu	-	-	-	+(32%)	-	+(43%)	+(67%)	+(31%)	-	+(22%)	-	-
Anti-Fas	-	-	+(48%)	+(5%)	-	-	-	-	+(9%)	-	-	+(4%)
H460-16-2	+(100%)	++	+++	++	++	+(33%)	+++	+++	++	+++	+++	+++

1 100 (+++) and in parenthesis, the percentage of cells stained. Representative histograms of
2 H460-16-2 antibodies were compiled for Figure 9 and evidence the binding characteristics,
3 inclusive of illustrated bimodal peaks in some cases. H460-16-2 bound 100 fold above
4 isotype control to a number of cancer cell types including melanoma and breast cancer
5 cells; 5 to 100 fold to lung, colon, prostate, and ovarian cancer cells. There was binding of
6 H460-16-2 antibodies to non-cancer cells, however that binding did not produce
7 cytotoxicity. This was evidence that binding was not necessarily predictive of the outcome
8 of antibody ligation of its cognate antigen, and was a non-obvious finding. This suggested
9 that the context of antibody ligation in different cells was determinative of cytotoxicity rather
10 than just antibody binding.

11

12 Example 5

13 **Normal Human Tissue Staining**

14 IHC studies were previously conducted to characterize H460-16-2 antigen
15 distribution in humans (S.N. 10/603,000). The current studies compare H460-16-2 to an
16 antibody directed against CD44 (L178) since the H460-16-2 antigen may be a cancer
17 variant of the standard form of CD44 as determined previously by biochemical methods.
18 Binding of antibodies to 59 normal human tissues was performed using a human, normal
19 organ tissue array (Imgenex, San Diego, CA). All primary antibodies (H460-16-2; L178
20 anti-CD44 (also known as anti-HCAM, BD PharMingen, Oakville, ON); and mouse IgG₁
21 negative control (Dako, Toronto, ON)) were diluted in antibody dilution buffer (Dako,
22 Toronto, ON) to a concentration of 5 µg/ml (found to be the optimal concentration in
23 previous optimization steps). The negative control antibody has been shown to be

1 negative to all mammalian tissues by the manufacturer. The procedure for IHC is as
2 follows.

3 Tissue sections were deparaffinized by drying in an oven at 58°C for 1 hour and
4 dewaxed by immersing in xylene 5 times for 4 minutes each in Coplin jars. Following
5 treatment through a series of graded ethanol washes (100%-75%) the sections were re-
6 hydrated in water. The slides were immersed in 10 mM citrate buffer at pH 6 (Dako,
7 Toronto, Ontario) then microwaved at high, medium, and low power settings for 5 minutes
8 each and finally immersed in cold PBS. Slides were then immersed in 3% hydrogen
9 peroxide solution for 6 minutes, washed with PBS three times for 5 minutes each, dried,
10 incubated with Universal blocking solution (Dako, Toronto, Ontario) for 5 minutes at room
11 temperature. H460-16-2, L178 or isotype control antibody (directed towards *Aspergillus*
12 *niger* glucose oxidase, an enzyme which is neither present nor inducible in mammalian
13 tissues) were diluted in antibody dilution buffer (Dako, Toronto, Ontario) to its working
14 concentration (5µg/mL for each antibody) and incubated for 1 hour at room temperature.
15 The slides were washed with PBS 3 times for 5 minutes each. Immunoreactivity of the
16 primary antibodies was detected/visualized with HRP conjugated secondary antibodies as
17 supplied (Dako Envision System, Toronto, Ontario) for 30 minutes at room temperature.
18 Following this step the slides were washed with PBS 3 times for 5 minutes each and a
19 color reaction developed by adding DAB (3,3'-diaminobenzidine tetrahydrachloride,
20 Dako, Toronto, Ontario) chromogen substrate solution for immunoperoxidase staining for
21 10 minutes at room temperature. Washing the slides in tap water terminated the
22 chromogenic reaction. Following counterstaining with Meyer's Hematoxylin (Sigma
23 Diagnostics, Oakville, ON), the slides were dehydrated with graded ethanols (75-100%)
24 and cleared with xylene. Using mounting media (Dako Faramount, Toronto, Ontario) the

slides were coverslipped. Slides were microscopically examined using an Axiovert 200 (Zeiss Canada, Toronto, ON) and digital images acquired and stored using Northern Eclipse Imaging Software (Mississauga, ON). Results were read, scored and interpreted by a pathologist.

Table 4 presents a summary of the results of H460-16-2 and L178 anti-CD44 staining of an array of normal human tissues. The staining of tissues with H460-16-2 is similar to that described previously (S.N. 10/603,000). It should be again noted that the antigen is generally not present on cells in the vital organs, including the liver, kidney, heart and lung. The H460-16-2 antibody does bind to macrophages and lymphocytes, and their presence is observed in some of the organs in these sections. However, there was a wider distribution and a higher intensity of staining of lymphocytes seen with the L178 anti-CD44 antibody (Figure 10).

Tissues that were positive for H460-16-2 were also usually positive for L178 anti-CD44 (sometimes to a greater intensity). Tissues that were negative for H460-16-2 were also generally negative for L178 anti-CD44 albeit there are a few exceptions such as one sample of liver (Figure 11) and esophagus. These results demonstrate that H460-16-2 binds to a slightly smaller subset of the tissues recognized by the L178 anti-CD44 antibody and within tissues the intensity of staining is also sometimes less. These results show that the antigen for H460-16-2 is not widely expressed on normal tissues, and that the antibody binds specifically to a limited number of tissues in humans. It also supports the biochemical data in that H460-16-2 is directed against an epitope of CD44, albeit to a slightly different variant than the one recognized by the L178 used for these IHC studies.

1
2
3
4
5
6
7
8

Table 4: Comparison of L178 anti-CD44 and H460-16-2 IHC on Human Normal Tissue

Abbreviations: *: Original pigmented stratum basale, **: Endogenous cytoplasmic pigment/back ground staining, ***: Stomach antrum (not stomach body), ****: Ovarian stroma only, SMF: Smooth muscle fibers, NR: The section is not representative, CS: The section is completely sloughed, PS: The section is partially sloughed.

Sec. No.	Organ	Anti-CD44	H460-16-2	Isotype Control
1	*Skin	+++ Keratinocytes of all epidermal layers, Fibroblasts, SMF of blood vessels	++ Keratinocytes of all epidermal layers, Fibroblasts	-
2	*Skin	+++ Keratinocytes of all epidermal layers, Fibroblasts	+++ Keratinocytes of all epidermal layers, Fibroblasts	-
3	Subcutis fat	-	-	-
4	Breast	+/- Ductular epithelium, Myoepithelium & Fibroblasts	+ Myoepithelium	-
5	Breast	+++ Myoepithelium & Fibroblasts	++ Myoepithelium & Fibroblasts	-
6	Spleen	+++ Lymphocytes predominantly in the periarteriolar area	++ Lymphocytes predominantly in the periarteriolar area	-
7	Spleen	+++ Lymphocytes predominantly in the periarteriolar area	+++ Lymphocytes predominantly in the periarteriolar area	-
8	Lymphnode	+++ Lymphocytes	+ Lymphocytes	-
9	Lymphnode	-	-	-
10	Skeletal muscle	+/- Blood vessels	+/- Blood vessels	-
11	Nasal Mucosa	- NR	CS	CS
12	Lung	+++ Macrophages & Fibroblasts	++ Macrophages	-
13	Lung	+++ Pneumocytes, Lymphocytes & Macrophages	+++ Lymphocytes & Macrophages	-
14	Bronchus	- NR	- NR	- NR
15	Heart	**	**	**
16	Salivary gland	+++ Acinar & Ductal epithelium, + Peripheral nerve	++ Acinar & ductal epithelium, + Peripheral nerve	-
17	Liver	+++ Kupffer cells	+++ Kupffer cells	-
18	Liver	+++ Kupffer cells	+++ Kupffer cells	-
19	Liver	++ Kupffer cells	-	-
20	Gall bladder	+++ Mucosal epithelium, Lymphocytes, Fibroblasts & SMF	+ Mucosal epithelium, +++ Lymphocytes	-
21	Pancreas	+ Acinar epithelium	+ Acinar epithelium	-
22	Pancreas	+++ Acinar epithelium	++ Acinar epithelium	-
23	Tonsil	+++ Lymphocytes all over the lymphatic nodules	++ Lymphocytes mainly in mantle zone	-
24	Esophagus	+ Squamous epithelium in basal layers, Fibroblasts	-	-
25	Esophagus	+++ Squamous epithelium in basal layers & Fibroblasts	+++ Squamous epithelium in basal layers & Fibroblasts	-
26	***Stomach body	+++ Gastric gland epithelium in basal parts of the glands & Lymphocytes	++ Gastric gland epithelium in basal parts of the glands, +++ Lymphocytes	-
27	***Stomach body	+++ Gastric gland epithelium in basal parts of the glands & Lymphocytes	+/- Gastric gland epithelium in basal parts of the glands, +++ Lymphocytes	-
28	Stomach antrum	+++ Lymphocytes & Fibroblasts	+++ Lymphocytes & Fibroblasts	-
29	Stomach Smooth muscle	++ Blood vessels & Fibroblasts	++ Blood vessels & Fibroblasts	-
30	Duodenum	+++ Glandular epithelium in deeper part of the glands & Lymphocytes	+/- Glandular epithelium in deeper part of the glands, ++ Lymphocytes	-
31	Small bowel	+/- Glandular epithelium in deeper part of the glands, +++ Lymphocytes in lamina propria	++ Lymphocytes in lamina propria	-
32	Small bowel	+++ Lymphocytes mainly in mantle zone of lymphatic nodules	+++ Lymphocytes mainly in mantle zone of lymphatic nodules	-
33	Appendix	++ Glandular epithelium, +++ Lymphocytes	+/- Mucosal epithelium, +++ Lymphocytes	-
34	Colon	+/- Glandular epithelium, +++ Lymphocytes, SMF & Ganglion cells	+++ Lymphocytes	-
35	Colon	+++ Lymphocytes, Ganglion cells & Endothelium of blood vessels	+++ Lymphocytes of lamina propria & Lymphatic nodules' mantle zone	-
36	Rectum	+++ Lymphocytes & Fibroblasts	+++ Lymphocytes & Fibroblasts	-
37	Kidney cortex	++ Endothelium of blood vessels & Interstitial Fibroblasts	+/- Interstitial fibroblasts	-
38	Kidney cortex	+/- Tubular epithelium & Interstitial fibroblasts	+/- Tubular epithelium & Interstitial fibroblasts	-
39	Kidney Medulla	+ SMF & Fibroblasts	+/- SMF & Fibroblasts	-
40	Urinary Bladder	+++ Superficial transitional epithelium, Lymphocytes & Macrophages	+/- Transitional epithelium, ++ Lymphocytes & Macrophages	-
41	Prostate	+++ Myoepithelium	+/- Glandular epithelium, +++ Myoepithelium	-
42	Prostate	+++ Myoepithelium & Fibroblasts	++ Myoepithelium	-
43	Seminal vesicle	+/- Mucosal epithelium, ++ Outer longitudinal SMF of muscularis externa	+/- Mucosal epithelium, ++ Outer longitudinal SMF of muscularis externa	-
44	Testis	+/- Interstitial blood vessels & Fibroblasts	+/- Interstitial fibroblasts	-
45	Endometrium proliferative	+++ Stromal fibroblasts	++ Stromal fibroblasts	-
46	Endometrium secretory	+++ Glandular epithelium & Stroma	++ Glandular epithelium & Stroma	-
47	Myometrium	+++ SMF & Fibroblasts	+++ SMF	-
48	Uterine cervix	+++ Squamous epithelium in deeper layers & Fibroblasts	+++ Squamous epithelium in deeper layers & Fibroblasts	-
49	Salpinx	+ Fibroblasts & SMF	+ Fibroblasts & SMF	-
50	***Ovary	+/- SMF of blood vessels	+/- SMF of blood vessels	-
51	Placenta, villi	++ Endothelium of blood vessels	+/- Endothelium of blood vessels	-
52	Placenta, villi	++ Endothelium of blood vessels	+/- Fibroblasts	-
53	Umbilical cord	-	-	-
54	Adrenal gland	+++/-	+++/-	+++/-
55	Thyroid	+/- Blood vessels & Para-follicular cells	+/- Blood vessels & Para-follicular cells	-
56	Thymus	+ Lymphocytes	+/- Lymphocytes	- PS
57	Brain white matter	-	-	-
58	Brain gray matter	-	-	-
59	Cerebellum	-	-	-

9

Example 6

Human Breast Tumor Tissue Staining

A previous IHC study was undertaken to determine the cancer association of the

H460-16-2 antigen with human breast cancers and whether the H460-16-2 antibody was

1 likely to recognize human cancers (S.N. 10/603,000). Currently, a comparison was made
2 for L178 anti-CD44 staining, and an antibody directed towards *Aspergillus niger* glucose
3 oxidase, an enzyme which is neither present nor inducible in mammalian tissues (negative
4 control). A breast cancer tissue array derived from 50 breast cancer patients and 9 samples
5 derived from non-neoplastic breast tissue in breast cancer patients was used (Imgenex
6 Corporation, San Diego, CA). The following information was provided for each patient:
7 age, sex, American Joint Committee on Cancer (AJCC) tumor stage, lymph node, estrogen
8 receptor (ER) and progesterone receptor (PR) status. The procedure for IHC from Example
9 5 was followed. All antibodies were used at a working concentration of 5 µg/ml.

10 Tables 5 and 6 provide summaries of H460-16-2 and L178 anti-CD44 antibody
11 staining of breast cancer tissue arrays respectively. Each array contained tumor samples
12 from 50 individual patients. Overall, 62 percent of the 50 patients tested were positive for
13 H460-16-2 antigen compared to 76 percent for CD44. In cases where both H460-16-2 and
14 L178 anti-CD44 stained the same tissue, 43% of the samples had higher intensity staining
15 with the L178 anti-CD44 in comparison to H460-16-2. For both the H460-16-2 and CD44
16 antigen, only 4 and 6 out of 10 normal breast tissue samples from breast cancer patients
17 were positive respectively. No clear correlation between estrogen and progesterone
18 receptor status was evident. It also did not appear to be a trend to greater positive
19 expression of the H460-16-2 and CD44 antigen with higher tumor stage.

20 Table 5: Human Breast Tumor IHC Summary for H460-16-2

			Binding Score						
		Total #	-	+/-	+	++	+++	Total Positive	% Positive
Patient Samples	Tumor	50	19	19	4	3	5	31	62
	Normal	10	0	1	0	2	1	4	40
ER Status	ER+	28	13	13	1	1	0	15	54
	ER-	22	6	8	3	0	5	16	73
	Unknown	0	0	0	0	0	0	0	0
PR Status	PR+	19	9	8	1	1	0	10	53
	PR-	30	8	14	3	0	5	22	73
	Unknown	1	0	1	0	0	0	1	100
AJCCTumorStage	T1	4	2	1	1	0	0	2	50
	T2	21	6	9	1	1	4	15	71
	T3	20	9	9	1	0	1	11	55
	T4	5	2	2	1	0	0	3	60

1

2 Table 6: Human Breast Tumor IHC Summary for Anti-CD44 (L178)

		Total #	-	+/-	+	++	+++	Total Positive	% Positive
Patient Samples	Tumor	50	12	20	3	7	8	39	78
	Normal	10	4	1	0	3	2	6	60
ER Status	ER+	28	7	14	1	4	2	21	75
	ER-	22	5	6	2	3	6	17	77
	Unknown	0	0	0	0	0	0	0	0
PR Status	PR+	19	5	9	1	2	2	14	74
	PR-	30	7	10	2	5	6	23	77
	Unknown	1	0	1	0	0	0	1	100
AJCCTumorStage	T1	4	1	2	0	0	1	3	75
	T2	21	5	6	2	2	6	16	76
	T3	20	6	9	1	3	1	14	70
	T4	5	0	3	0	2	0	5	100

3

4 The H460-16-2 staining was specific for cancerous cells in comparison to normal
5 cells as demonstrated in Figure 12 where stromal cells were clearly negative and sheets of
6 malignant cells were highly positive. The cellular localization pattern seen with the H460-
7 16-2 antigen was confined to the cell membrane in the majority of cases. The L178 CD44
8 antibody stained more breast cancer samples and showed a higher degree of membrane
9 than cytoplasmic localization compared to H460-16-2 (Table 7). L178 anti-CD44 also
10 stained malignant cells of Paget's disease, which was not the case for H460-16-2 (Figure
11 13). L178 anti-CD44 stained the same samples of normal tissue from breast cancer patients
12 as H460-16-2 (plus 1 that was partially sloughed for the H460-16-2 sample). These results
13 suggest the antigen for H460-16-2 may be expressed by almost two thirds of breast cancer

1 patients. The staining pattern showed that in patient samples, the antibody is highly
2 specific for malignant cells and the H460-16-2 antigen is localized to the cell membrane
3 thereby making it an attractive drugable target. The similar albeit more limited staining of
4 H460-16-2 versus L178 anti-CD44 antibody again demonstrates the likelihood of the
5 H460-16-2 epitope being a more restricted variant of CD44.

6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

Sec. No.	Sex	Age	Diagnosis	Anti-CD44	H460-16-2	Isotype Control
1	F	28	Infiltrating ductal carcinoma	+/-	+/-	-
2	F	71	Solid papillary carcinoma	++ M	+/-	-
3	F	26	Infiltrating ductal carcinoma	+/-	-	-
4	F	43	Infiltrating ductal carcinoma	+/-	-	-
5	F	39	Infiltrating ductal carcinoma	+/-	+/- Tumor & Necrotic area	-
6	F	46	Ductal carcinoma <i>in situ</i>	+/-	-	-
7	F	47	Infiltrating ductal carcinoma	+++ M	+++ M	-
8	M	67	Infiltrating ductal carcinoma	-Tumor, + Stroma only	-	-
9	F	33	Infiltrating ductal carcinoma	-Tumor, ++ Stroma only	- Tumor, ++ Stroma	-
10	F	47	Infiltrating ductal carcinoma	+/-	+/-	-
11	F	49	Invasive lobular carcinoma	- Tumor, ++ Infiltrating lymphocytes	-	-
12	F	46	Infiltrating ductal carcinoma	++ M	+/-	-
13	F	39	Infiltrating ductal carcinoma	+/-	-	-
14	F	43	Infiltrating lobular carcinoma	+ MC	+/-	-
15	F	54	Infiltrating lobular carcinoma	-Tumor, + Stroma	+/-	-
16	F	58	Infiltrating ductal carcinoma	++ M Tumor, +++ Stroma	+ MCTumor, ++ Necrotic area	-
17	F	37	Infiltrating ductal carcinoma	- Tumor, + Stroma	+/-	-
18	F	43	Infiltrating ductal carcinoma	++ M Tumor, +++ Stroma	+ MCTumor, +++ Stroma	-
19	F	51	Infiltrating ductal carcinoma	+++ M	++ M	-
20	F	80	Medullary carcinoma	+++ M	+++ MC	-
21	F	36	Infiltrating ductal carcinoma	+++ M	+++ M	-
22	F	59	Infiltrating ductal carcinoma	+/- Tumor, ++ Stroma	+/- Tumor, ++ Stroma	-
23	F	34	Ductal carcinoma <i>in situ</i>	+/-	+/- Tumor & Necrotic area	-
24	F	54	Infiltrating ductal carcinoma	+/-	+/-	-
25	F	47	Infiltrating ductal carcinoma	+++ M	+/-	-
26	F	53	Infiltrating ductal carcinoma	+Tumor, +++ Stroma	- Tumor, ++ Stroma	-
27	F	59	Infiltrating ductal carcinoma	+/- Tumor, +++ Lymphocytes	+/- Tumor, +++ Lymphocytes	-
28	F	60	Signet ring cell carcinoma	- Tumor, +/- Stroma	-	-
29	F	37	Infiltrating ductal carcinoma	+/-	+/-	-
30	F	46	Infiltrating ductal carcinoma	++ Tumor, +++ Infiltrating Lymphocytes	+/- Tumor & Stroma	-
31	F	35	Infiltrating ductal carcinoma	+/-	-	-
32	F	47	Infiltrating ductal carcinoma	- Tumor, ++ Necrotic cells	- Tumor, +/- Necrotic cells	-
33	F	54	Infiltrating ductal carcinoma	+/- Tumor & Stroma	-	-
34	F	47	Infiltrating ductal carcinoma	+++ M	+++ M	-
35	F	41	Infiltrating ductal carcinoma	-	-	-
36	F	38	Infiltrating ductal carcinoma	++ M	+/-	-
37	F	55	Infiltrating ductal carcinoma	-Tumor, +/- Stroma	-	-
38	F	65	Infiltrating ductal carcinoma	- Tumor, ++ Stroma	+/- Stroma	-
39	M	66	Infiltrating ductal carcinoma	-	-	-
40	F	44	Infiltrating ductal carcinoma	+/-	- Tumor, + Infiltrating lymphocytes	-
41	F	52	Metastatic carcinoma in lymph node	+/- Tumor, +++ Infiltrating lymphocytes	+/- Tumor & Stroma	-
42	F	32	Metastatic carcinoma in lymph node	+/-	-	-
43	F	58	Metastatic carcinoma in lymph node	++ M	+ MC	-
44	F	52	Metastatic carcinoma in lymph node	-	-	-
45	F	58	Metastatic carcinoma in lymph node	+/-	+/-	-
46	F	38	Metastatic carcinoma in lymph node	+/- Tumor, +++ Infiltrating lymphocytes	-Tumor, + Lymphocytes	-
47	F	45	Metastatic carcinoma in lymph node	+++ M	+ MC	-
48	F	45	Metastatic carcinoma in lymph node	+ M	+/-	-
49	F	29	Metastatic carcinoma in lymph node	+++ M	+++ M	-
50	F	61	Metastatic carcinoma in lymph node	+/- Tumor, ++ Stroma	+/- Tumor, ++ Stroma	-
*51	F	46	Nipple	+++ Keratinocytes	++ Keratinocytes	-
*52	F	47	Nipple	+/- Tumor cells	-	-
*53	F	40	Normal breast	-	-	-
*54	F	43	Normal breast	++ Myoepithelium	+++ Myoepithelium	-
*55	F	40	Normal breast	++ Myoepithelium	++ Myoepithelium	-
*56	F	40	Normal breast	+++ Myoepithelium & Fibroblasts	+/- Myoepithelium & Fibroblasts	-
*57	F	45	Normal breast	-	-	-
*58	F	44	Normal breast	-	-	-
*59	F	37	Normal breast	-	-	-
60	F	51	Normal breast	++ Myoepithelium & Fibroblasts	- PS	-

Table 7: Comparison of L178 anti-CD44 and H460-16-2 IHC on Human Tumor and Normal Breast Tissue

Abbreviations: *: Non-neoplastic breast tissue in breast cancer patient, PS: The section is partially sloughed, M: Membrane staining, C: Cytoplasmic staining.

Example 7

Human Tumor Tissue Staining

To determine whether the H460-16-2 antigen is expressed on other human cancer tissues in addition to breast cancer, H460-16-2 was previously used on a multiple human tumor tissue array (S.N. 10/603,000; Imgenex, San Diego, CA). In furthering those studies, the staining pattern of H460-16-2 was compared to that of L178 anti-CD44 The

1 following information was provided for each patient: age, sex, organ and diagnosis. The
2 staining procedure used was the same as the one outlined in Example 5. The same
3 negative control antibody was used as described for the human breast tumor tissue array.
4 All antibodies were used at a working concentration of 5 µg/mL.

5 As outlined in Table 8, H460-16-2 stained a number of various human cancers
6 besides breast with results consistent with those described previously (S.N. 10/603,000).
7 As seen with the breast cancers, H460-16-2 staining was localized on the membrane
8 (Figure 14) and in some of the cancers also within the cytoplasm of cancerous cells. L178
9 anti-CD44 antibody had greater membrane versus cytoplasmic staining with again a higher
10 percentage of tumor tissues staining positive and with greater intensity than that observed
11 with H460-16-2 (Figure 15).

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

Sec. No.	Age	Sex	Organ	Diagnosis	Anti-CD44	H460-16-2	Isotype Control
1	59	M	Skin	Malignant Melanoma	+++ M	+++ M	-
2	25	F	Skin	SSC	+/	-	-
3	50	F	Breast	Infiltrating ductal carcinoma	+++ Tumor & Stroma	+ Tumor, +++ Stroma	-
4	57	F	Breast	Invasive papillary carcinoma	+/ Tumor & Stroma	+/ Tumor & Stroma	-
5	35	F	Breast	Infiltrating lobular carcinoma	+ M	+/	-
6	40	M	Lymph node	Malignant lymphoma, Immunoplastic	+++ M	+++ M	-
7	58	M	Lymph node	Metastatic adenocarcinoma from stomach	+++ M	+/	-
8	53	F	Bone	Osteosarcoma	+++ M	+ M/C	-
9	26	M	Bone	Giant cell tumor	++ M	+ M/C	-
10	40	M	bone	Chondro-sarcoma	CS	CS	CS
11	51	F	Soft tissue	Liposarcoma	-	-	-
12	47	F	Soft tissue	Neuro-fibromatosis	++ M/C	+ M/C	-
13	74	M	Nasal cavity	Inverted papilloma	+++ M	++ M	-
14	57	M	Larynx	SCC	+++ M	+++ M	-
15	60	M	Lung	Adenocarcinoma	+++ M	+/	-
16	51	F	Lung	SCC	+++ M	+++ M/C	-
17	68	F	Lung	Adenocarcinoma	++ M	+/	-
18	60	M	Lung	Small cell carcinoma	+/ Necrotic area	+/	-
19	88	F	Tongue	SCC	+++ M	+++ M	.*
20	34	F	Parotid gland	Pleomorphic adenoma	++ M/C	-	-
21	50	F	Parotid gland	Warthin tumor	+++ M/C	+++ M/C	-
22	40	F	Parotid gland	Pleomorphic adenoma	+++ M	++ M/C	-
23	56	M	Submandibular gland	Salivary duct carcinoma	+/	-	-
24	69	F	Liver	Cholangio-carcinoma	+/ Tumor, ++ Stroma	+/	-
25	51	M	Liver	Metastatic gastric carcinoma	- Tumor, +++ Necrotic area	-	-
26	64	M	Liver	HCC	+/ Tumor, ++ Extracellular secretion	+/ Tumor, ++ Extracellular secretion	-
27	62	F	Gall bladder	Adenocarcinoma	+++ Tumor	++ Tumor & Lymphocytes	-
28	64	F	Pancreas	Adenocarcinoma	++ M	++ M/C	-
29	68	M	Esophagus	SCC	++ M	+/	-
30	73	M	Stomach	Adenocarcinoma, Poorly differentiated	+/ Tumor, ++ Stroma	+ M/C Tumor & Stroma	-
31	63	M	Stomach	Adenocarcinoma, Moderately differentiated	+/ Tumor, +++ Infiltrating lymphocytes	++ M/C	-
32	59	F	Stomach	Signet ring cell carcinoma	+++ M	++ M/C	-
33	62	M	Stomach	Malignant lymphoma	+++ M	+++ M/C	.*
34	51	M	Stomach	Borderline stromal tumor	+/ Stroma	-	-
35	42	M	Small intestine	Malignant stromal tumor	++ M	-	-
36	52	F	Appendix	Pseudomyxomatous peritonitis	PS, +++ Infiltrating lymphocytes	-	-
37	53	M	Colon	Adenocarcinoma	+ M/C Tumor & Stroma	+ M/C	-
38	67	M	Rectum	Adenocarcinoma	+/	++ M	-
39	75	F	Kidney	Transitional cell carcinoma	++ M	+ M/C	-
40	54	F	Kidney	Renal cell carcinoma	+++ M	+/	-
41	75	F	Kidney	Renal cell carcinoma	+/	+/	-
42	65	M	Urinary bladder	Poorly differentiated carcinoma	+++ Infiltrating lymphocytes	++ M/C	-
43	67	M	Urinary bladder	Transitional cell carcinoma, High grade	+/ Stroma	-	-
44	62	M	Prostate	Adenocarcinoma	+++ M	+++ M	.*
45	30	M	Testis	Seminoma	+/ Tumor, +++ Stroma	+/	-
46	68	F	Uterus	Endometrial adenocarcinoma	++ Stroma mainly	++ Stroma mainly	-
47	57	F	uterus	Leiomyosarcoma	++ C	+ PS	-
48	45	F	uterus	Leiomyoma	++ C	+ C	-
49	63	F	Uterine cervix	SCC	+++ M	+++ M	-
50	12	F	Ovary	Endodermal sinus tumor	+/	-	-
51	33	F	Ovary	Mucinous adenocarcinoma	+ M/C	-	-
52	70	F	Ovary	Fibrosarcoma	-	-	-
53	67	F	Adrenal gland	Cortical carcinoma	-	-	.*
54	61	F	Adrenal gland	Pheochromocytoma	-	-	-
55	54	M	Thyroid	Papillary carcinoma	++ M	++ M/C	-
56	38	F	Thyroid	Minimally invasive follicular carcinoma	+++ M	++ M	-
57	74	M	Thyroid	Thyroidoma	++ M/C	+/	-
58	66	F	Brain	Meningioma	+/	-	-
59	62	M	Brain	Glioblastoma multiforme	+++ M	+++ M	-

Table 8: Comparison of L178 anti-CD44 and H460-16-2 IHC on Multiple Human Tumor Types
Abbreviations: M: Membrane staining, C: Cytoplasmic staining, *: Background stain, CS: The section is completely sloughed, PS: The section is partially sloughed, F: The section is folded, SSC: Squamous cell carcinoma, HCC: Hepatocellular carcinoma.

Therefore, it appears that the H460-16-2 antigen is not solely found on the membranes of breast cancers but also on the membrane of a large variety of tumor types. These results indicate that H460-16-2 has potential as a therapeutic drug in a wide variety of tumor types besides breast. Again, the similar yet distinct staining pattern of H460-16-2 compared to L178 anti-CD44 implies that H460-16-2 is recognizing an epitope present on a variant of CD44.

The preponderance of evidence shows that H460-16-2 mediate anti-cancer effects through ligation of a carbohydrate dependent conformational epitope present on a variant

1 of CD44. It has been shown, in example 3, H460-16-2 antibody can be used to
2 immunoprecipitate the cognate antigen from expressing cells such as MB-231 cells.
3 Further it could be shown that the H460-16-2 antibody could be used in detection of cells
4 and/or tissues which express a CD44 antigenic moiety which specifically binds thereto,
5 utilizing techniques illustrated by, but not limited to FACS, cell ELISA or IHC.

6 Thus, it could be shown that the immunoprecipitated H460-16-2 antigen can inhibit
7 the binding of H460-16-2 to such cells or tissues using such FACS, cell ELISA or IHC
8 assays. Further, as with the H460-16-2 antibody, other anti-CD44 antibodies could be used
9 to immunoprecipitate and isolate other forms of CD44 antigen, and the antigen can also be
10 used to inhibit the binding of those antibodies to the cells or tissues that express the antigen
11 using the same types of assays. It could also be shown that if an anti-CD44 antibody that
12 recognizes all forms of CD44 (i.e. a pan-CD44 antibody) were used to isolate its cognate
13 antigen, then that antigen could also inhibit the binding of H460-16-2 antigen to cells or
14 tissues that express that antigen, thus also demonstrating the binding of H460-16-2 to an
15 epitope of CD44 on cells and tissues expressing that antigen. Alternatively, a comparison
16 of H460-16-2 and pan-CD44 antibody in assays such as competitive binding assays.
17 ELISA, cell ELISA, FACS or the like, where both antibodies are present can also
18 demonstrate the binding of H460-16-2 to an epitope of CD44 on cells and tissues
19 expressing that antigen.

20 All patents and publications mentioned in this specification are indicative of the
21 levels of those skilled in the art to which the invention pertains. All patents and
22 publications are herein incorporated by reference to the same extent as if each individual
23 publication was specifically and individually indicated to be incorporated by reference.

1 It is to be understood that while a certain form of the invention is illustrated, it is
2 not to be limited to the specific form or arrangement of parts herein described and shown.
3 It will be apparent to those skilled in the art that various changes may be made without
4 departing from the scope of the invention and the invention is not to be considered limited
5 to what is shown and described in the specification. One skilled in the art will readily
6 appreciate that the present invention is well adapted to carry out the objects and obtain the
7 ends and advantages mentioned, as well as those inherent therein. Any oligonucleotides,
8 peptides, polypeptides, biologically related compounds, methods, procedures and
9 techniques described herein are presently representative of the preferred embodiments, are
10 intended to be exemplary and are not intended as limitations on the scope. Changes therein
11 and other uses will occur to those skilled in the art which are encompassed within the spirit
12 of the invention and are defined by the scope of the appended claims. Although the
13 invention has been described in connection with specific preferred embodiments, it should
14 be understood that the invention as claimed should not be unduly limited to such specific
15 embodiments. Indeed, various modifications of the described modes for carrying out the
16 invention which are obvious to those skilled in the art are intended to be within the scope
17 of the following claims.

18

19